

FOUR POSITIVE FORMULAE FOR TYPE A QUIVER POLYNOMIALS

ALLEN KNUTSON, EZRA MILLER, AND MARK SHIMOZONO

ABSTRACT. We give four positive formulae for the (equioriented type A) quiver polynomials of Buch and Fulton [BF99]. All four formulae are combinatorial, in the sense that they are expressed in terms of combinatorial objects of certain types: Zelevinsky permutations, lacing diagrams, Young tableaux, and pipe dreams (also known as rc-graphs). Three of our formulae are multiplicity-free and geometric, meaning that their summands have coefficient 1 and correspond bijectively to components of a torus-invariant scheme. The remaining (presently non-geometric) formula is a variant of the conjecture of Buch and Fulton in terms of factor sequences of Young tableaux [BF99]; our proof of it proceeds by way of a new characterization of the tableaux counted by quiver constants. All four formulae come naturally in “doubled” versions, two for *double quiver polynomials*, and the other two for their stable limits, the double quiver functions, where setting half the variables equal to the other half specializes to the ordinary case.

Our method begins by identifying quiver polynomials as multidegrees [BB82, Jos84, BB85, Ros89] via equivariant Chow groups [EG98]. Then we make use of Zelevinsky’s map from quiver loci to open subvarieties of Schubert varieties in partial flag manifolds [Zel85]. Interpreted in equivariant cohomology, this lets us write double quiver polynomials as ratios of double Schubert polynomials [LS82] associated to Zelevinsky permutations; this is our first formula. In the process, we provide a simple argument that Zelevinsky maps are scheme-theoretic isomorphisms (originally proved in [LM98]). Writing double Schubert polynomials in terms of pipe dreams [FK96] then provides another geometric formula for double quiver polynomials, via [KM03a]. The combinatorics of pipe dreams for Zelevinsky permutations implies an expression for limits of double quiver polynomials in terms of products of Stanley symmetric functions [Sta84]. A degeneration of quiver loci (orbit closures of GL on quiver representations) to unions of products of matrix Schubert varieties [Ful92, KM03a] identifies the summands in our Stanley function formula combinatorially, as lacing diagrams that we construct based on the strands of Abeasis and Del Fra in the representation theory of quivers [AD80]. Finally, we apply the combinatorial theory of key polynomials to pass from our lacing diagram formula to a double Schur function formula in terms of peelable tableaux [RS95a, RS98], and from there to our formula of Buch–Fulton type.

CONTENTS

Introduction	3
Overview	3
Acknowledgments	4
The four formulae	4
Proofs via double versions and limits	11
Related notions and extensions	12
A note on the field \mathbb{k}	13

Date: 16 January 2006.

AK was partly supported by the Sloan Foundation and NSF.

EM and MS were partly supported by the NSF.

Section 1. Geometry of quiver loci	14
1.1. Quiver loci and ideals	14
1.2. Rank, rectangle, and lace arrays	15
1.3. The Zelevinsky map	18
1.4. Quiver polynomials as multidegrees	21
Section 2. Double quiver polynomials	24
2.1. Double Schubert polynomials	24
2.2. Double quiver polynomials	25
2.3. Ratio formula for quiver polynomials	27
Section 3. Lacing diagrams	30
3.1. Geometry of lacing diagrams	30
3.2. Minimum length lacing diagrams	32
Section 4. Degeneration of quiver loci	34
4.1. Orbit degenerations	34
4.2. Quiver degenerations	35
4.3. Multidegrees of quiver degenerations	36
4.4. Rank stability of components	37
Section 5. Pipe dreams for Zelevinsky permutations	40
5.1. Pipe dream formula for double quiver polynomials	40
5.2. Pipes to laces	43
5.3. Rank stability of pipe dreams	44
Section 6. Component formulae	47
6.1. Double Stanley symmetric functions	49
6.2. Double quiver functions	51
6.3. All components are lacing diagram orbit closures	54
6.4. Stable double component formula	55
Section 7. Quiver constants	57
7.1. Demazure characters	57
7.2. Schubert polynomials as sums of Demazure characters	59
7.3. Quiver constants are Stanley coefficients	61
7.4. Peelable tableaux	63
Section 8. Factor sequences from peelable tableaux	65
8.1. Factor sequences	65
8.2. Zelevinsky peelable tableaux	68
8.3. Bijection to factor sequences	69
8.4. The Buch–Fulton conjecture	73
References	76

REFERENCES

- [AD80] S. Abeasis and A. Del Fra, *Degenerations for the representations of an equioriented quiver of type A_m* , Boll. Un. Mat. Ital. Suppl. (1980) no. 2, 157–171.
- [ADK81] S. Abeasis, A. Del Fra, and H. Kraft, *The geometry of representations of A_m* , Math. Ann. **256** (1981), no. 3, 401–418.
- [BB93] Nantel Bergeron and Sara Billey, *RC-graphs and Schubert polynomials*, Experiment. Math. **2** (1993), no. 4, 257–269.
- [BJS93] Sara C. Billey, William Jockusch, and Richard P. Stanley, *Some combinatorial properties of Schubert polynomials*, J. Algebraic Combin. **2** (1993), no. 4, 345–374.
- [BB82] W. Borho and J.-L. Brylinski, *Differential operators on homogeneous spaces. I. Irreducibility of the associated variety for annihilators of induced modules*, Invent. Math. **69** (1982), no. 3, 437–476.
- [BB85] W. Borho and J.-L. Brylinski, *Differential operators on homogeneous spaces. III. Characteristic varieties of Harish–Chandra modules and of primitive ideals*, Invent. Math. **80** (1985), no. 1, 1–68.
- [Bri97] Michel Brion, *Equivariant Chow groups for torus actions*, Transform. Groups **2** (1997), no. 3, 225–267.
- [BF99] Anders Skovsted Buch and William Fulton, *Chern class formulas for quiver varieties*, Invent. Math. **135** (1999), no. 3, 665–687.
- [BFR03] Anders S. Buch, László M. Fehér and Richárd Rimányi, *Positivity of quiver coefficients through Thom polynomials*, to appear in Adv. Math.
- [BKTY02] Anders S. Buch, Andrew Kresch, Harry Tamvakis, and Alexander Yong, *Schubert polynomials and quiver formulas*, Duke Math. J. **122** (2004), no. 1, 125–143.
- [Buc01a] Anders Skovsted Buch, *Stanley symmetric functions and quiver varieties*, J. Algebra **235** (2001), no. 1, 243–260.
- [Buc01b] Anders Skovsted Buch, *On a conjectured formula for quiver varieties*, J. Algebraic Combin. **13** (2001), no. 2, 151–172.
- [Buc02] Anders Skovsted Buch, *Grothendieck classes of quiver varieties*, Duke Math. J. **115** (2002), no. 1, 75–103.
- [Buc03] Anders Skovsted Buch, *Alternating signs of quiver coefficients*, J. Amer. Math. Soc. **18** (2005), no. 1, 217–237.
- [Dem74] M. Demazure, *Une nouvelle formule des caractères*, Bull. Sci. Math. (2) **98** (1974), no. 3, 163–172.
- [EG87] Paul Edelman and Curtis Greene, *Balanced tableaux*, Adv. in Math. **63** (1987), no. 1, 42–99.
- [EG98] Dan Edidin and William Graham, *Equivariant intersection theory*, Invent. Math. **131** (1998), no. 3, 595–634.
- [Eis95] David Eisenbud, *Commutative algebra, with a view toward algebraic geometry*, Graduate Texts in Mathematics, vol. 150, Springer-Verlag, New York, 1995.
- [FR02a] László Fehér and Richárd Rimányi, *Schur and Schubert polynomials as Thom polynomials—cohomology of moduli spaces*, Cent. Eur. J. Math. **1** (2003), no. 4, 418–434.
- [FR02b] László Fehér and Richárd Rimányi, *Classes of degeneracy loci for quivers: the Thom polynomial point of view*, Duke Math. J. **114** (2002), no. 2, 193–213.
- [FK94] Sergey Fomin and Anatol N. Kirillov, *Grothendieck polynomials and the Yang–Baxter equation*, 1994, Proceedings of the Sixth Conference in Formal Power Series and Algebraic Combinatorics, DIMACS, pp. 183–190.
- [FK96] Sergey Fomin and Anatol N. Kirillov, *The Yang–Baxter equation, symmetric functions, and Schubert polynomials*, Proceedings of the 5th Conference on Formal Power Series and Algebraic Combinatorics (Florence, 1993), Discrete Math. **153** (1996), no. 1-3, 123–143.
- [FS94] Sergey Fomin and Richard P. Stanley, *Schubert polynomials and the nil-Coxeter algebra*, Adv. Math. **103** (1994), no. 2, 196–207.
- [Ful92] William Fulton, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*, Duke Math. J. **65** (1992), no. 3, 381–420.
- [Ful97] William Fulton, *Young tableaux, with applications to representation theory and geometry*, London Mathematical Society Student Texts **35**, Cambridge University Press, Cambridge, 1997.
- [Ful98] William Fulton, *Intersection theory*, second ed., Springer-Verlag, Berlin, 1998.

- [FP98] William Fulton and Piotr Pragacz, *Schubert varieties and degeneracy loci*, Appendix J by the authors in collaboration with I. Ciocan-Fontanine. Lecture Notes in Mathematics 1689, Springer-Verlag, Berlin, 1998.
- [Ful99] William Fulton, *Universal Schubert polynomials*, Duke Math. J. **96** (1999), no. 3, 575–594.
- [Gia04] G. Z. Giambelli, *Ordine di una varietà più ampia di quella rappresentata coll'annullare tutti i minori di dato ordine estratti da una data matrice generica di forme*, Mem. R. Ist. Lombardo 3rd series **11** (1904) 101–135.
- [Hai92] Mark Haiman, *Dual equivalence with applications, including a conjecture of Proctor*, Discrete Math. **99** (1992), no. 1–3, 79–113.
- [Jos84] Anthony Joseph, *On the variety of a highest weight module*, J. Algebra **88** (1984), no. 1, 238–278.
- [KS95] Michael Kalkbrener and Bernd Sturmfels, *Initial complexes of prime ideals*, Adv. Math. **116** (1995), no. 2, 365–376.
- [Kaz97] M. É. Kazarian, *Characteristic classes of singularity theory*, The Arnold-Gelfand mathematical seminars, 325–340, Birkhäuser Boston, Boston, MA, 1997.
- [KM03a] Allen Knutson and Ezra Miller, *Gröbner geometry of Schubert polynomials*, to appear in Ann. of Math (2).
- [KM03b] Allen Knutson and Ezra Miller, *Subword complexes in Coxeter groups*, Adv. Math. **184** (2004), no. 1, 161–176.
- [Kog00] Mikhail Kogan, *Schubert geometry of flag varieties and Gel'fand-Cetlin theory*, Ph.D. thesis, Massachusetts Institute of Technology, 2000.
- [Las90] Alain Lascoux, *Anneau de Grothendieck de la variété de drapeaux*, The Grothendieck Festschrift, Vol. III, 1–34, Progr. Math., **88**, Birkhäuser Boston, Boston, MA, 1990.
- [Las01] Alain Lascoux, *Transition on Grothendieck polynomials*, Physics and combinatorics, 2000 (Nagoya), 164–179, World Sci. Publishing, River Edge, NJ, 2001.
- [Las03] Alain Lascoux, *Double crystal graphs*, Studies in Memory of Issai Schur (Chevaleret/Rehovot, 2000), 95–114, Progr. Math., **210**, Birkhäuser Boston, Boston, MA, 2003.
- [Len02] Cristian Lenart, *A unified approach to combinatorial formulas for Schubert polynomials*, J. Algebraic Combin. **20** (2004), no. 3, 263–299.
- [LM98] V. Lakshmibai and Peter Magyar, *Degeneracy schemes, quiver schemes, and Schubert varieties*, Internat. Math. Res. Notices (1998), no. 12, 627–640.
- [LS82] Alain Lascoux and Marcel-Paul Schützenberger, *Structure de Hopf de l'anneau de cohomologie et de l'anneau de Grothendieck d'une variété de drapeaux*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), no. 11, 629–633.
- [LS85] Alain Lascoux and Marcel-Paul Schützenberger, *Schubert polynomials and the Littlewood-Richardson rule*, Lett. Math. Phys. **10** (1985), no. 2-3, 111–124.
- [LS89] Alain Lascoux and Marcel-Paul Schützenberger, *Noncommutative Schubert polynomials*, Func. Anal. Appl. **23** (1989), no. 3, 223–225 (1990).
- [LS90] Alain Lascoux and Marcel-Paul Schützenberger, *Keys and standard bases*, Invariant theory and tableaux (Minneapolis, MN, 1988), 125–144, IMA Vol. Math. Appl., 19, Springer, New York, 1990.
- [Lus90] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), no. 2, 447–498.
- [Mac91] I. G. Macdonald, *Notes on Schubert polynomials*, Publ. LACIM **6**, UQAM, Montréal, 1991.
- [Mag98] Peter Magyar, *Schubert polynomials and Bott-Samelson varieties*, Comment. Math. Helv. **73** (1998), no. 4, 603–636.
- [Mil03] Ezra Miller, *Alternating formulae for K-theoretic quiver polynomials*, Duke Math. J. **128** (2005), no. 1, 1–17.
- [MS04] Ezra Miller and Bernd Sturmfels, *Combinatorial commutative algebra*, Graduate Texts in Mathematics, **227**, Springer-Verlag, New York, 2005.
- [Ram85] A. Ramanathan, *Schubert varieties are arithmetically Cohen-Macaulay*, Invent. Math. **80** (1985), no. 2, 283–294.
- [RR85] S. Ramanan and A. Ramanathan, *Projective normality of flag varieties and Schubert varieties*, Invent. Math. **79** (1985), no. 2, 217–224.
- [Ros89] W. Rossmann, *Equivariant multiplicities on complex varieties*, Orbites unipotentes et représentations, III, Astérisque no. 173–174, (1989), 11, 313–330.

- [RS95a] Victor Reiner and Mark Shimozono, *Plactification*, J. Algebraic Combin. **4** (1995), no. 4, 331–351.
- [RS95b] Victor Reiner and Mark Shimozono, *Key polynomials and a flagged Littlewood-Richardson rule*, J. Combin. Theory Ser. A **70** (1995), no. 1, 107–143.
- [RS98] Victor Reiner and Mark Shimozono, *Percentage-avoiding, northwest shapes and peelable tableaux*, J. Combin. Theory Ser. A **82** (1998), no. 1, 1–73.
- [Sta84] R. P. Stanley, *On the number of reduced decompositions of elements of Coxeter groups*, European J. Combin. **5** (1984), no. 4, 359–372.
- [Tho55] R. Thom, *Les singularités des applications différentiables*, Ann. Inst. Fourier, Grenoble **6** (1955–1956), 43–87.
- [Yon03] Alexander Yong, *On combinatorics of quiver component formulas*, J. Algebraic Combin. **21** (2005) 351–371.
- [Zel85] A. V. Zelevinskii, *Two remarks on graded nilpotent classes*, Uspekhi Mat. Nauk **40** (1985), no. 1(241), 199–200.

MATHEMATICS DEPARTMENT, UC BERKELEY, BERKELEY, CALIFORNIA
E-mail address: `allenk@math.berkeley.edu`

MATHEMATICAL SCIENCES RESEARCH INSTITUTE, BERKELEY, CALIFORNIA
E-mail address: `ezra@math.umn.edu`

MATHEMATICS DEPARTMENT, VIRGINIA TECH, BLACKSBURG, VIRGINIA
E-mail address: `mshimo@math.vt.edu`