



FROM KINETIC TO FLUID MODELS OF LIQUID CRYSTALS BY THE MOMENT METHOD

PIERRE DEGOND*

Institut de Mathématiques de Toulouse; UMR5219
Université de Toulouse; CNRS UPS
F-31062 Toulouse Cedex 9, France

AMIC FROUVELLE

CEREMADE, CNRS, Université Paris-Dauphine
Université PSL, 75016 Paris, France
and
CNRS, Université de Poitiers, UMR 7348
Laboratoire de Mathématiques et Applications (LMA)
86000 Poitiers, France

JIAN-GUO LIU

Department of Physics and Department of Mathematics
Duke University
Durham, NC 27708, USA

In memory of Bob Glassey

ABSTRACT. This paper deals with the convergence of the Doi-Navier-Stokes model of liquid crystals to the Ericksen-Leslie model in the limit of the Deborah number tending to zero. While the literature has investigated this problem by means of the Hilbert expansion method, we develop the moment method, i.e. a method that exploits conservation relations obeyed by the collision operator. These are non-classical conservation relations which are associated with a new concept, that of Generalized Collision Invariant (GCI). In this paper, we develop the GCI concept and relate it to geometrical and analytical structures of the collision operator. Then, the derivation of the limit model using the GCI is performed in an arbitrary number of spatial dimensions and with non-constant and non-uniform polymer density. This non-uniformity generates new terms in the Ericksen-Leslie model.

2020 *Mathematics Subject Classification.* 35Q35, 76A10, 76A15, 82C22, 82C70, 82D30, 82D60.

Key words and phrases. Doi equation, Maier-Saupe interaction, Q-tensor, Deborah number, Ericksen-Leslie equations, Oseen-Franck energy, order parameter, generalized collision invariant.

PD is a visiting professor of the Department of Mathematics, Imperial College London, UK.

AF acknowledges support from the Project EFI ANR-17-CE40-0030 of the French National Research Agency.

JGL acknowledges support from the Department of Mathematics, Imperial College London, under Nelder Fellowship award and the National Science Foundation under Grants DMS-1812573 and DMS-2106988.

* Corresponding author: Pierre Degond.

1. Introduction. We consider the Doi kinetic model of liquid crystals coupled with the Navier-Stokes equation for the fluid solvent. We investigate the limit of the Deborah number tending to zero by means of a moment method. The limit model is a system of fluid equations named the Ericksen-Leslie model [25, 37, 59]. In classical kinetic theory, there are two methods to derive fluid equations, the Hilbert expansion method [7, 11, 26, 34] and the moment method [5, 52]. However, for a number of kinetic models including the Doi model, only the Hilbert method can be used. Indeed, the moment method is subject to a condition on the number of conservation relations satisfied by the collision operator and this condition is not satisfied by the Doi model. This is why the Hilbert expansion method is the only method developed in the literature so far (see e.g. [25, 37, 59]). In the present work, we address the question whether the moment method can be used for the Doi kinetic model.

The moment method relies on the requirement that the space of collision invariants (the quantities conserved by the collisions operator) has the same dimensions as the number of free parameters in the local equilibrium distribution function. This requirement is not satisfied for the Doi model. The goal of this paper is to show that the moment method can still be used for this model but necessitates a weaker concept of collision invariant, that of “generalized collision invariant” or GCI. This concept has first been introduced in [23] to derive the fluid dynamic limit of the Vicsek model [55]. Since then, the GCI concept has been applied to a variety of collective dynamics models [17, 18, 19, 21, 22, 30]. The present work is its first application to visco-elastic fluid models.

Visco-elastic fluids have been the subject of an abundant literature (see e.g. [2, 3, 16, 24, 32, 58] for reviews). The Doi model is one of the most fundamental models of visco-elastic fluids [24]. It models an assembly of polymer molecules flowing in an incompressible fluid (the solvent). The polymer molecules are assumed to be rigid spheroids mutually interacting through alignment and subject to noise. They are represented by a distribution function of their position and orientation. After Onsager and Maier-Saupe [50, 53], alignment accounts for the volume exclusion interaction between the molecules. Alignment is of nematic type, i.e. invariant if the head and tail of the molecules are flipped. Following Landau and de Gennes [16], the interaction depends on the Q-tensor, a quadratic quantity which respects this invariance. The fluid solvent is modelled by the incompressible Navier-Stokes equations. Polymer molecules are transported by the fluid and rotated by the fluid gradients. In turn, they influence the fluid through extra-stresses depending on their distribution function. The mathematical theory of this model has been investigated in [45, 54, 62] and for active particles, in [13].

The Doi model involves a dimensionless parameter, the Deborah number which describes the alignment rate of the polymer molecules. When this parameter goes to zero, the distribution of orientations tends to a profile which depends on two parameters, the polymer density ρ and the polymer molecules average orientation Ω which are functions of space and time. In the case of a constant density ρ , it is shown in [25, 37, 59] that the mean orientation satisfies a transport-diffusion equation. Its coupling with the Navier-Stokes equations leads to the so-called Ericksen-Leslie system [27, 38]. The convergence is formal in [25, 37] and rigorous in [59]. In all cases, the method relies on the Hilbert expansion. There is an abundant mathematical literature on the Ericksen-Leslie system per se [35, 42, 43, 44, 60].

Here, our goal is to provide a formal convergence proof of the Doi model to the Ericksen-Leslie model using the moment method. Specifically, we will derive the appropriate GCI concept, discuss its rationale and relation to the Hilbert expansion method. There are several motivations to develop a moment method even if a Hilbert expansion theory already exists. The first one is that the GCI concept has an underlying geometrical structure which is worth highlighting and may lead to new structural properties of the Doi model. The second reason is that a mathematical theory based on the moment method often requires less regularity than the Hilbert expansion method (compare e.g. [4] with [7]). This potentially opens the way to simpler convergence proofs from the Doi to the Ericksen-Leslie models. The third reason is that the moment method naturally leads to the development of efficient numerical methods [33, 39] which might enable us to handle the complexity of the Doi kinetic model in a systematic way.

Aside to this main goal, we will pursue two secondary goals. The first one is to provide a treatment of the small Deborah number limit in arbitrary dimension. So far, this has only been done in dimension 3. This extension is made possible by Wang and Hoffman [56] who have determined the spatially uniform equilibria in any dimension. Although dimension three is the physically relevant case, there are several reasons for considering an arbitrary dimension. The first one is that the use of dimension 3 often conceals simple structures under dimension-specific concepts and notations. Then, as argued in [12], the use of an arbitrary dimension often reveals hidden and interesting mathematical properties. Finally, fluid-dynamic equations are based on simple postulates that may be adapted to other objects such as information flow in abstract spaces of large dimensions.

The second side goal is to investigate the effect of a spatially non-uniform density of polymer molecules. To the best of our knowledge, earlier work on the small Deborah number limit [25, 37, 59] have assumed the density of polymer molecules to be constant. Investigation of Ericksen-Leslie models with non-uniform order parameter has been made in the literature [8, 9, 27, 40, 41, 46], but none has explicitly linked this non-uniform order parameter to the non-uniform polymer density (as it should as we will see) and derived these models from kinetic theory. Non-uniform polymer density results in modifications of the equations for the mean director Ω and for the extra-stresses that will be highlighted in this work.

The organization of this paper is as follows: Section 2 gives an exposition of the Doi-Navier-Stokes model and the small Deborah number scaling. Section 3 is devoted to the statement of the main result, namely the formal convergence of the Doi-Navier-Stokes model to the Ericksen-Leslie model in the zero Deborah number limit. Section 4 describes the local equilibria. Section 5 develops the GCI concept for the Doi model and discusses it. In Section 6, the limiting equations of the Doi model when the Deborah number tends to zero are derived. Conclusions and perspectives are drawn in Section 7. Auxiliary results stated in Sections 2, 3, 5 and 6 are proved in appendices A, B, C and D respectively.

2. Kinetic model for rod-like polymer suspensions and scaling.

2.1. The Doi equation. In this paper, we consider the Doi model [16, 24, 25, 37, 54, 57, 59], where polymer molecules are identified as spheroids. We consider the semi-dilute regime [24, 25, 57] where a volume-exclusion interaction potential needs to be incorporated. We neglect the inertia of the polymer molecules. Following [25, 37, 57], we describe the polymer molecules by a kinetic distribution function

$f(x, \omega, t)$ where $x \in \mathbb{R}^n$ is the position, $\omega \in \mathbb{S}^{n-1}$ is the molecule orientation and $t \geq 0$ is the time. We let \mathbb{S}^{n-1} be the unit $(n-1)$ -dimensional sphere and since ω and $-\omega$ refer to the same molecular orientation, we impose

$$f(x, \omega, t) = f(x, -\omega, t). \quad (1)$$

Let $u(x, t) \in \mathbb{R}^n$ be the fluid velocity. In general, the dimension $n = 2$ or 3 but the theory will be developed for any value of n . The equation for f (the so-called Doi equation) reads as follows:

$$\partial_t f + \nabla_x \cdot (uf) + \nabla_\omega \cdot (f(\Lambda P_{\omega^\perp} E - W)\omega) = D \nabla_\omega \cdot (\nabla_\omega f + \frac{1}{k_B T} f \nabla_\omega U_f^R). \quad (2)$$

Here, D denotes the rotational diffusivity, T , the fluid temperature and k_B , the Boltzmann constant. The tensors E and W are respectively the symmetric and anti-symmetric parts of the velocity gradient, given by

$$E = \frac{1}{2}(\nabla_x u + \nabla_x u^T), \quad W = \frac{1}{2}(\nabla_x u - \nabla_x u^T). \quad (3)$$

The symbols ∇_x and $\nabla_x \cdot$ refer to the spatial gradient and divergence operators while ∇_ω , $\nabla_\omega \cdot$ to the gradient and divergence operators on the sphere \mathbb{S}^{n-1} respectively. The notation $\nabla_x u$ refers to the gradient tensor of u defined by $(\nabla_x u)_{ij} = \partial_{x_i} u_j$ and the exponent ‘T’ indicates the transpose. The dimensionless quantity Λ is related to the aspect ratio (ratio between the semi-axes) of the spheroidal polymer molecules. Finally, $P_{\omega^\perp} = \text{Id} - \omega \otimes \omega$ for $\omega \in \mathbb{S}^{n-1}$ denotes the projection operator of vectors onto the normal hyperplane to ω . Throughout this paper, Id denotes the identity matrix and if $u = (u_i)_{i=1, \dots, n}$ and $v = (v_i)_{i=1, \dots, n}$ are two vectors, $u \otimes v$ denotes their tensor product, i.e. the $n \times n$ tensor $(u \otimes v)_{ij} = u_i v_j$. For two $n \times n$ tensors S and S' , SS' stands for the matrix product of S and S' , hence the meaning of $P_{\omega^\perp} E$. The surface measure on the sphere will be normalized, meaning that $\int_{\mathbb{S}^{n-1}} d\omega = 1$.

The quantity U_f^R is the interaction potential stemming from volume exclusion between the polymer molecules. In the Maier-Saupe theory [50], this interaction potential reads

$$U_f^R(x, \omega, t) = k_B T \nu \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} \frac{1}{R^n} K\left(\frac{|x-x'|}{R}\right) (1 - (\omega \cdot \omega')^2) f(x', \omega', t) d\omega' dx', \quad (4)$$

where ν is the potential strength. Following the formalism proposed by [25, 57], a spatial non-locality is introduced by means of the kernel $K: [0, \infty) \rightarrow [0, \infty)$, $\xi \mapsto K(\xi)$ which describes the influence of two neighboring molecules. Specifically, two molecules separated by a distance ξ influence each other with strength $\frac{1}{R^n} K(\frac{\xi}{R})$, where R is the typical interaction range. The kernel K satisfies $\int_{\mathbb{R}^n} K(|x|) dx = 1$. An equivalent expression of U_f^R is

$$U_f^R(x, \omega, t) = k_B T \nu \rho_f^R \left[-(\omega \cdot Q_f^R \omega) + \frac{n-1}{n} \right], \quad (5)$$

where ρ_f^R and Q_f^R are the locally averaged particle density and orientational de Gennes Q-tensor given by

$$\rho_f^R(x, t) = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} \frac{1}{R^n} K\left(\frac{|x-x'|}{R}\right) f(x', \omega, t) d\omega dx', \quad (6)$$

$$(\rho_f^R Q_f^R)(x, t) = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} \frac{1}{R^n} K\left(\frac{|x-x'|}{R}\right) \left(\omega \otimes \omega - \frac{1}{n} \text{Id}\right) f(x', \omega, t) d\omega dx'. \quad (7)$$

Note that Q_f^R is a trace-free symmetric matrix obtained by averaging $\omega \otimes \omega - \frac{1}{n} \text{Id}$ over the probability distribution $\rho_f^R(x, t)^{-1} R^{-n} K(|x - x'|/R) f(x', \omega, t) d\omega dx'$. Consequently, thanks to the min-max theorem, its eigenvalues λ satisfy the inequality

$$-\frac{1}{n} \leq \lambda \leq 1 - \frac{1}{n}. \tag{8}$$

The following fully local versions of the polymer density and orientational tensor:

$$\rho_f = \int_{\mathbb{S}^{n-1}} f d\omega = \lim_{R \rightarrow 0} \rho_f^R, \tag{9}$$

$$\rho_f Q_f = \int_{\mathbb{S}^{n-1}} \left(\omega \otimes \omega - \frac{1}{n} \text{Id} \right) f d\omega = \lim_{R \rightarrow 0} \rho_f^R Q_f^R, \tag{10}$$

will also be useful. From (5), it follows that

$$\frac{1}{k_B T} \nabla_\omega U_f^R(x, \omega, t) = -2\nu \rho_f^R P_{\omega^\perp} Q_f^R \omega,$$

so that an alternate formulation of the Doi equation (2) is given by

$$\begin{aligned} \partial_t f + \nabla_x \cdot (uf) + \nabla_\omega \cdot (f(\Lambda P_{\omega^\perp} E - W)\omega) \\ = D(\Delta_\omega f - 2\nu \rho_f^R \nabla_\omega \cdot (f P_{\omega^\perp} Q_f^R \omega)). \end{aligned} \tag{11}$$

We note that Eq. (11) preserves the symmetry constraint (1). The second and third term at the left-hand side of (11) model passive transport of the polymer molecules by the fluid: the second term corresponds to translation of the molecules by the fluid velocity and the third term to their rotation by the gradient of the fluid velocity. Here, we assume that the polymer molecules can be described by spheroids, i.e. ellipsoids, in which $n - 1$ semi-axes b are equal. The aspect ratio p is the ratio a/b where a is the remaining semi-axis. The quantity Λ is related to p by $\Lambda = \frac{p^2 - 1}{p^2 + 1}$. In particular, $\Lambda \in [-1, 1]$ and $\Lambda = 1$ for infinitely thin rods, $\Lambda = 0$ for spheres, and $\Lambda = -1$ for infinitely flat disks. The rotation operator is derived from Jeffery's equation [36]. The first term at the right-hand side of (11) describes Brownian effects due to rotational diffusion. We neglect translational diffusivity, as it is usually much smaller than rotational diffusivity [16]. The second term at the right-hand side of (11) takes into account the volume exclusion interaction between the molecules and drives the distribution to that of a system of fully aligned polymer molecules. To measure the degree of alignment of the molecules, one introduces

$$\chi_f = \frac{n}{n - 1} \lambda_f \text{ with } \lambda_f = \text{the largest eigenvalue of } Q_f, \tag{12}$$

where Q_f is given by (10). This quantity can be seen as the order parameter for the distribution f . We have $\chi_f \in (0, 1)$. If f is close to the uniform distribution on the sphere, which corresponds to a fully disordered distribution of polymer orientations, then χ_f is close to 0. By contrast, if f is close to $\frac{1}{2}(\delta_\Omega + \delta_{-\Omega})$ where Ω is any vector on \mathbb{S}^{n-1} , which corresponds to a fully aligned distribution of polymer orientations in the direction $\pm\Omega$, then, χ_f is close to 1.

To ensure thermodynamic consistency, one introduces the polymer free energy [25]:

$$\mathcal{A}^R(t) = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} [k_B T (f \log f - f) + \frac{1}{2} U_f^R f] dx d\omega.$$

From (4), it is easy to check that the quantity $\int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} U_f^R g dx d\omega$ defined for two functions f and g of (x, ω) is a symmetric bilinear form. Then the functional

derivative $\mu_f^R = \frac{\delta \mathcal{A}^R}{\delta f}$, also referred to as the chemical potential, is given by

$$\mu_f^R = k_B T \log f + U_f^R = k_B T \left(\log f - \nu \rho_f^R \left[(\omega \cdot Q_f^R \omega) - \frac{n-1}{n} \right] \right). \quad (13)$$

Thus,

$$\nabla_\omega \mu_f^R = k_B T \left(\frac{\nabla_\omega f}{f} - 2\nu \rho_f^R P_{\omega^\perp} Q_f^R \omega \right), \quad (14)$$

so that (2) can also be written:

$$\partial_t f + \nabla_x \cdot (u f) + \nabla_\omega \cdot (f (\Lambda P_{\omega^\perp} E - W) \omega) = \frac{D}{k_B T} \nabla_\omega \cdot (f \nabla_\omega \mu_f^R). \quad (15)$$

The right-hand side of (15) can be viewed as describing the steepest descent in the direction of the minimum of the polymer free energy. This is also known as the maximal dissipation principle. Using Green's formula, we have the following identity (provided f vanishes fast enough at infinity), whose proof is sketched in Appendix A.1:

$$\frac{d\mathcal{A}^R}{dt} = \int_{\mathbb{R}^n} \sigma_f^R : \nabla_x u \, dx - \int_{\mathbb{R}^n} F_f^R \cdot u \, dx - \frac{D}{k_B T} \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} f |\nabla_\omega \mu_f^R|^2 \, dx \, d\omega, \quad (16)$$

where σ_f^R is the extra-stress tensor and F_f^R is a body force, given by :

$$\sigma_f^R = \int_{\mathbb{S}^{n-1}} \left(\Lambda (\omega \otimes \nabla_\omega \mu_f^R)_s + (\omega \otimes \nabla_\omega \mu_f^R)_a \right) f \, d\omega, \quad F_f^R = - \int_{\mathbb{S}^{n-1}} \nabla_x \mu_f^R f \, d\omega. \quad (17)$$

Here, for two $n \times n$ tensors $S = (S_{ij})_{ij=1,\dots,n}$ and $S' = (S'_{ij})_{ij=1,\dots,n}$, we denote by $S : S' = S_{ij} S'_{ij}$ their contraction (with the repeated index summation convention) while S_s and S_a are respectively the symmetric and antisymmetric parts of S namely $S_s = \frac{1}{2}(S + S^T)$, $S_a = \frac{1}{2}(S - S^T)$. Contractions and tensor products will be defined and noted similarly for tensors of higher order.

2.2. The Navier-Stokes equations. The Doi equation (2) (or equivalently, (11) or (15)) is coupled to the Navier-Stokes equation for the fluid velocity, which is written [24, 25, 57]:

$$\rho_F (\partial_t u + u \cdot \nabla_x u) + \nabla_x p = \nabla_x \cdot (\sigma_f^R + \tau_u + \mathcal{T}_{f,u}) + F_f^R, \quad (18)$$

$$\nabla_x \cdot u = 0. \quad (19)$$

Here ρ_F is the fluid mass density. The extra-stress tensor σ_f^R is given by (17) while τ_u and $\mathcal{T}_{f,u}$ are contributions of the fluid and polymer molecules to the viscous stresses respectively given by

$$\tau_u = 2\eta E, \quad \mathcal{T}_{f,u} = \zeta \frac{k_B T}{D} \rho_f \mathbb{T}_f : E,$$

with the fourth order orientational tensor \mathbb{T}_f given by

$$\rho_f \mathbb{T}_f = \int_{\mathbb{S}^{n-1}} \omega^{\otimes 4} f \, d\omega. \quad (20)$$

For a $n \times n$ tensor S , its divergence $\nabla_x \cdot S$ denotes the vector defined by $(\nabla_x \cdot S)_j = \partial_{x_i} S_{ij}$ (using the repeated index summation convention). As above, $\mathbb{T}_f : E$ denotes the contraction of \mathbb{T}_f and E with respect to two indices. Although \mathbb{T}_f is a fourth order tensor, it is symmetric, so which pair of its indices is concerned by the contraction is indifferent. The quantity η is the fluid viscosity. Using the divergence-free condition (19), we remark that $\nabla_x \cdot \tau_u = \eta \Delta_x u$. The quantity ζ is a dimensionless number. In [24], for the dilute polymer regime in dimension 3, it is

shown that $\zeta = \frac{1}{2}$. But this derivation requires the use of the Oseen tensor which has dimensional dependence [12] and thus, the value of ζ changes with the dimension. Moreover, even in dimension 3, in the semi-dilute regime considered here, the value of ζ may be different from $\frac{1}{2}$ [24, Section 9.5.1]. So, we shall consider ζ as a free parameter of the model.

We have the following expression for the extra-stress:

$$\sigma_f^R = nk_B T \Lambda \rho_f Q_f + \int_{\mathbb{S}^{n-1}} \left[\frac{\Lambda + 1}{2} \omega \otimes \nabla_\omega U_f^R + \frac{\Lambda - 1}{2} \nabla_\omega U_f^R \otimes \omega \right] f d\omega. \quad (21)$$

However, although more complicated, the following expression, which is valid if f is a solution of the Doi equation (2), will turn out to be more useful:

$$\begin{aligned} \sigma_f^R = & \frac{k_B T \Lambda}{D} \frac{1}{2} \rho_f [\Lambda(EQ_f + Q_f E) + Q_f W - WQ_f + \frac{2\Lambda}{n} E - 2\Lambda \mathbb{T}_f : E \\ & - D_t Q_f] + \frac{1}{2} \int_{\mathbb{S}^{n-1}} (\omega \otimes \nabla_\omega U_f^R - \nabla_\omega U_f^R \otimes \omega) f d\omega, \end{aligned} \quad (22)$$

where

$$D_t = \partial_t + u \cdot \nabla_x, \quad (23)$$

is the material derivative. Eq. (21) results from the first equation of (17) after insertion of (14). Eq. (22) is obtained by multiplying Doi's equation (15) by $\omega \otimes \omega - \frac{1}{n} \text{Id}$ and integrating with respect to ω , followed by some algebra. These computations have been done in [25, 37, 59] for $n = 3$ and are sketched in Appendix A.2 for any n .

The rationale for involving σ_f^R and F_f^R in the coupling between the Navier-Stokes equations (18) and the Doi equation (2) is thermodynamical consistency. Indeed, we have the following total free energy dissipation identity (provided spatial boundary terms vanish in the integrations by parts):

$$\frac{d}{dt} \mathcal{E}^R + \mathcal{D}^R = 0, \quad (24)$$

where \mathcal{E}^R is the total free energy (sum of the fluid and polymer free energies):

$$\mathcal{E}^R(t) = \int_{\mathbb{R}^n} \frac{1}{2} \rho_F |u|^2 dx + \mathcal{A}^R,$$

and \mathcal{D}^R is the total free energy dissipation:

$$\mathcal{D}^R(t) = \frac{D}{k_B T} \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} f |\nabla_\omega \mu_f^R|^2 dx d\omega + \int_{\mathbb{R}^n} (2\eta E : E + \frac{k_B T \zeta}{D} \rho_f \mathbb{T}_f : (E \otimes E)) dx,$$

where now, $\mathbb{T}_f : (E \otimes E)$ indicates the contraction of the fourth order tensors \mathbb{T}_f and $E \otimes E$ with respect to all four indices. We have omitted the dependence of E on u for simplicity.

2.3. Scaling. We now introduce a suitable scaling of this model. Let x_0 , t_0 and ρ_0 be space, time and polymer density units and let $u_0 = x_0/t_0$, $f_0 = \rho_0$, $\sigma_0 = k_B T \rho_0$, $p_0 = \rho_F u_0^2$, $F_0 = \sigma_0/x_0$, $U_0 = k_B T$ be units for velocity, distribution function, stress tensor, fluid pressure, elastic force and potential respectively. Then, we introduce the following dimensionless quantities:

$$\text{De} = \frac{1}{Dt_0}, \quad \text{Re} = \frac{u_0 x_0 \rho_F}{\eta}, \quad \text{Er} = \frac{\eta D}{k_B T \rho_0}, \quad \alpha = \nu \rho_0, \quad \bar{R} = \frac{R}{x_0}.$$

The dimensionless quantities De, Re and Er are the classical Deborah, Reynolds and Ericksen numbers, which respectively encode the relaxation time of the polymer molecular assembly to equilibrium, the ratio of inertial to viscous forces in the fluid

and the ratio between the viscous and extra stresses. The parameters α and \bar{R} are measures of the molecular interaction intensity and range respectively. The other dimensionless parameters of the model are ζ and Λ . Introducing scaled variables $x' = x/x_0$, $t' = t/t_0$ and unknowns $f(x, \omega, t) dx d\omega = \rho_0 f'(x', \omega, t') dx' d\omega$, $u(x, t) = u_0 u'(x', t')$, \dots , we can deduce the following dimensionless form of the Doi model (dropping the primes for clarity):

$$\partial_t f + \nabla_x \cdot (uf) + \nabla_\omega \cdot (f(\Lambda P_{\omega^\perp} E - W)\omega) = \frac{1}{\text{De}} \nabla_\omega \cdot (\nabla_\omega f + f \nabla_\omega U_f^{\bar{R}}), \quad (25)$$

with

$$U_f^{\bar{R}} = \alpha \rho_f^{\bar{R}} \left[-(\omega \cdot Q_f^{\bar{R}} \omega) + \frac{n-1}{n} \right],$$

and $\rho_f^{\bar{R}}$, $Q_f^{\bar{R}}$ given by (6), (7) with R replaced by \bar{R} . The polymer free energy is now given by

$$\mathcal{A}^{\bar{R}}(t) = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} (f \log f - f + \frac{1}{2} U_f^{\bar{R}} f) dx d\omega$$

and the chemical potential $\mu_f^{\bar{R}} = \frac{\delta \mathcal{A}^{\bar{R}}}{\delta f}$ by

$$\mu_f^{\bar{R}} = \log f + U_f^{\bar{R}} = \log f - \alpha (\omega \cdot \rho_f^{\bar{R}} Q_f^{\bar{R}} \omega) + \alpha \frac{n-1}{n} \rho_f^{\bar{R}}.$$

Thus, the expression at the right-hand side of (25) is equivalently written

$$\nabla_\omega \cdot (\nabla_\omega f + f \nabla_\omega U_f^{\bar{R}}) = \nabla_\omega \cdot (f \nabla_\omega \mu_f^{\bar{R}}) = \Delta_\omega f - 2\alpha \rho_f^{\bar{R}} \nabla_\omega (f P_{\omega^\perp} Q_f^{\bar{R}} \omega).$$

The scaled Navier-Stokes equation reads as follows

$$\begin{aligned} \partial_t u + u \cdot \nabla_x u + \nabla_x p &= \frac{1}{\text{Re}} \nabla_x \cdot (\tau_u + \frac{1}{\text{Er}} \mathcal{T}_{f,u}) + \frac{1}{\text{Re Er De}} (\nabla_x \cdot \sigma_f^{\bar{R}} + F_f^{\bar{R}}), \\ \nabla_x \cdot u &= 0, \\ \tau_u &= 2E, \quad \mathcal{T}_{f,u} = \zeta \rho_f \mathbb{T}_f : E, \end{aligned}$$

with $\sigma_f^{\bar{R}}$, $F_f^{\bar{R}}$ given by (17) with R replaced by \bar{R} and ρ_f , \mathbb{T}_f given by (9), (20). Expressions (21), (22) for the stress tensor are scaled into

$$\begin{aligned} \sigma_f^{\bar{R}} &= n\Lambda \rho_f Q_f + \int_{\mathbb{S}^{n-1}} \left[\frac{\Lambda+1}{2} \omega \otimes \nabla_\omega U_f^{\bar{R}} + \frac{\Lambda-1}{2} \nabla_\omega U_f^{\bar{R}} \otimes \omega \right] f d\omega. \\ &= \text{De} \frac{\Lambda}{2} \rho_f [\Lambda(EQ_f + Q_f E) + Q_f W - WQ_f + \frac{2\Lambda}{n} E - 2\Lambda \mathbb{T}_f : E \\ &\quad - D_t Q_f] + \frac{1}{2} \int_{\mathbb{S}^{n-1}} (\omega \otimes \nabla_\omega U_f^{\bar{R}} - \nabla_\omega U_f^{\bar{R}} \otimes \omega) f d\omega, \quad (26) \end{aligned}$$

with Q_f still given by (10). The free-energy dissipation identity is still written as (24) with $\mathcal{E}^{\bar{R}}$ and $\mathcal{D}^{\bar{R}}$ now given by

$$\begin{aligned} \mathcal{E}^{\bar{R}} &= \frac{1}{2} \int_{\mathbb{R}^n} |u|^2 dx + \frac{1}{\text{Re Er De}} \mathcal{A}^{\bar{R}}, \\ \mathcal{D}^{\bar{R}} &= \frac{1}{\text{Re}} \int_{\mathbb{R}^n} |\nabla_x u|^2 dx + \frac{1}{\text{Re Er}} \zeta \int_{\mathbb{R}^n} \rho_f \mathbb{T}_f : (E \otimes E) dx \\ &\quad + \frac{1}{\text{Re Er De}^2} \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} f |\nabla_\omega \mu_f^{\bar{R}}|^2 dx d\omega, \quad (27) \end{aligned}$$

where, for a $n \times n$ tensor S , $|S|$ denotes the Frobenius norm of S , i.e. $|S|^2 = \text{Tr}\{S^T S\}$.

The goal of this article is to investigate the limit of the Deborah number De tending to zero through the use of the new “generalized collision invariant” concept. In doing so, we will keep the parameters Re , Er and α of order unity. As for \bar{R} , following [25, 59], we make the scaling $\bar{R} = \mathcal{O}(\sqrt{De})$. This scaling assumption is analogous to the weakly non-local interaction scaling of the Vicsek model [20]. As we may choose the time and space units independently, we assume:

$$De = \varepsilon, \quad \bar{R} = \sqrt{\varepsilon}, \quad \varepsilon \rightarrow 0,$$

and assume Re , Er and α independent of ε . A straightforward Taylor expansion shows that

$$\rho_f^{\sqrt{\varepsilon}} = \rho_f + \varepsilon \beta \Delta_x \rho_f + \mathcal{O}(\varepsilon^2), \quad \rho_f^{\sqrt{\varepsilon}} Q_f^{\sqrt{\varepsilon}} = \rho_f Q_f + \varepsilon \beta \Delta_x (\rho_f Q_f) + \mathcal{O}(\varepsilon^2),$$

where

$$\beta = \frac{1}{2n} \int_{\mathbb{R}^n} K(|x|) |x|^2 dx. \tag{28}$$

Then, we can expand $U_f^{\sqrt{\varepsilon}} = U_f^0 + \varepsilon U_f^1 + \mathcal{O}(\varepsilon^2)$, $\mu_f^{\sqrt{\varepsilon}} = \mu_f^0 + \varepsilon \mu_f^1 + \mathcal{O}(\varepsilon^2)$ with

$$U_f^0 = \alpha \rho_f \left[-(\omega \cdot Q_f \omega) + \frac{n-1}{n} \right], \quad \mu_f^0 = \log f + U_f^0 \tag{29}$$

$$U_f^1 = \mu_f^1 = \beta \Delta_x U_f^0. \tag{30}$$

Straightforward computations show that

$$\int_{\mathbb{S}^{n-1}} \omega \otimes \nabla_\omega U_f^0 f d\omega = -2\alpha \rho_f^2 [Q_f^2 + \frac{1}{n} Q_f - \mathbb{T}_f : Q_f], \tag{31}$$

so that the left-hand side of (31) is a symmetric tensor. We deduce that the integral term in (26) is $\mathcal{O}(\varepsilon)$, so that $\sigma_f^{\sqrt{\varepsilon}} = \mathcal{O}(\varepsilon)$. Additionally, similar computations as for (31) lead to

$$\int_{\mathbb{S}^{n-1}} (\omega \otimes \nabla_\omega U_f^1 - \nabla_\omega U_f^1 \otimes \omega) f d\omega = 2\alpha \beta \rho_f [\Delta_x (\rho_f Q_f) Q_f - Q_f \Delta_x (\rho_f Q_f)].$$

So, we can write $\sigma_f^{\bar{R}} = \varepsilon \sigma_f^1 + \mathcal{O}(\varepsilon^2)$ with

$$\begin{aligned} \sigma_f^1 = & \rho_f \frac{\Lambda}{2} [\Lambda (E Q_f + Q_f E) + Q_f W - W Q_f + \frac{2\Lambda}{n} E - 2\Lambda \mathbb{T}_f : E \\ & - D_t Q_f] + \alpha \beta \rho_f [\Delta_x (\rho_f Q_f) Q_f - Q_f \Delta_x (\rho_f Q_f)]. \end{aligned} \tag{32}$$

We also note that $F_f^{\sqrt{\varepsilon}} = -\nabla_x \varphi_f^0 + \varepsilon F_f^1 + \mathcal{O}(\varepsilon^2)$, with

$$\varphi_f^0 = \rho_f - \frac{\alpha}{2} \rho_f^2 [Q_f : Q_f - \frac{n-1}{n}], \quad F_f^1 = - \int_{\mathbb{S}^{n-1}} \nabla_x \mu_f^1 f d\omega. \tag{33}$$

We let $\tilde{p}^\varepsilon = p^\varepsilon + \frac{1}{\varepsilon} \frac{1}{ReEr} \varphi_f^0$. We will omit the tilde below for simplicity. Since the $\mathcal{O}(\varepsilon^2)$ terms in all these developments have no contribution to the limit model when $\varepsilon \rightarrow 0$ (at the leading order), we will just ignore them.

We finally get the following perturbation problem:

$$\begin{aligned} \partial_t f^\varepsilon + \nabla_x \cdot (u^\varepsilon f^\varepsilon) + \nabla_\omega \cdot (f^\varepsilon (\Lambda P_{\omega^\perp} E^\varepsilon - W^\varepsilon) \omega) \\ + 2\alpha \beta \nabla_\omega \cdot (f^\varepsilon P_{\omega^\perp} \Delta_x (\rho_{f^\varepsilon} Q_{f^\varepsilon}) \omega) = \frac{1}{\varepsilon} \left(\Delta_\omega f^\varepsilon - 2\alpha \rho_{f^\varepsilon} \nabla_\omega \cdot (f^\varepsilon P_{\omega^\perp} Q_{f^\varepsilon} \omega) \right), \end{aligned} \quad (34)$$

$$\begin{aligned} \partial_t u^\varepsilon + u^\varepsilon \cdot \nabla_x u^\varepsilon + \nabla p^\varepsilon = \frac{1}{\operatorname{Re}} \left\{ \Delta_x u^\varepsilon + \right. \\ \left. \frac{1}{\operatorname{Er}} \left[\zeta \nabla_x \cdot (\rho_{f^\varepsilon} \mathbb{T}_{f^\varepsilon} : E^\varepsilon) + \nabla_x \cdot \sigma_{f^\varepsilon}^1 + F_{f^\varepsilon}^1 \right] \right\}, \end{aligned} \quad (35)$$

$$\nabla_x \cdot u^\varepsilon = 0, \quad (36)$$

where $\sigma_{f^\varepsilon}^1$ is given by (32) and $F_{f^\varepsilon}^1$ by (33).

We define the transport operator $T_u(f)$ (for a given time-dependent vector field $u: \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}^n$) and the collision operator $C(f)$ by

$$\begin{aligned} T_u(f) &= \partial_t f + \nabla_x \cdot (u f) + \nabla_\omega \cdot (f (\Lambda P_{\omega^\perp} E - W) \omega) \\ &\quad + 2\alpha \beta \nabla_\omega \cdot (f P_{\omega^\perp} \Delta_x (\rho_f Q_f) \omega), \end{aligned} \quad (37)$$

$$C(f) = \Delta_\omega f - 2\alpha \rho_f \nabla_\omega \cdot (f P_{\omega^\perp} Q_f \omega) = \nabla_\omega \cdot (f \nabla_\omega \mu_f^0) \quad (38)$$

$$= \nabla_\omega \cdot (\nabla_\omega f + f \nabla_\omega U_f^0), \quad (39)$$

so that (34) is written

$$T_{u^\varepsilon}(f^\varepsilon) = \frac{1}{\varepsilon} C(f^\varepsilon). \quad (40)$$

We note that $\mu_f^0 = \frac{\delta \mathcal{A}^0}{\delta f}$ is the functional derivative of the free energy $\mathcal{A}^0 = \lim_{\varepsilon \rightarrow 0} \mathcal{A}^{\sqrt{\varepsilon}}$ given by

$$\mathcal{A}^0(t) = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} (f \log f - f + \frac{1}{2} U_f^0 f) dx d\omega, \quad (41)$$

and recall that U_f^0 and μ_f^0 are given by (29). We refer to [25] for the formulation of the free energy dissipation identity for the whole model (34)-(36).

3. Main result.

3.1. Preliminaries. The purpose of this paper is to derive the limit of model (34)-(36) when $\varepsilon \rightarrow 0$. Before stating the result, we need a few preliminaries. We note that C given by (38) operates on the variable ω only and leaves (x, t) as parameters. This justifies the definition:

Definition 3.1. A function $f: \mathbb{S}^{n-1} \rightarrow \mathbb{R}$, $\omega \mapsto f(\omega)$ is called an equilibrium of C if and only if it satisfies

$$C(f) = 0. \quad (42)$$

Remark 1. We note that f is an equilibrium if and only if f is a critical point of the free energy functional \mathcal{A}^0 given by (41) in the spatially homogeneous case (i.e. when f is a function of ω only and integration with respect to x in the definition of \mathcal{A}^0 is ignored) [1, 47, 48, 59]. Moreover, such equilibria will be called “stable” if they correspond to local minimizers of this free energy (see [28] for $n = 2$, [1, 29, 47, 48] for $n = 3$ and [31] for $n = 4$).

The equilibria will attract the dynamics as $\varepsilon \rightarrow 0$ and their determination is of key importance. For this purpose, we introduce the Gibbs distributions:

Definition 3.2 (Gibbs distribution). Let S be a trace-free symmetric matrix. Then, the Gibbs distribution G_S associated with S is given by:

$$G_S(\omega) = \frac{1}{Z_S} e^{\omega \cdot S \omega}, \quad Z_S = \int_{\mathbb{S}^{n-1}} e^{\omega \cdot S \omega} d\omega. \quad (43)$$

Next, we introduce the

Definition 3.3 (Normalized prolate uniaxial trace-free tensor). Let $\Omega \in \mathbb{P}^{n-1} := \mathbb{S}^{n-1}/\{\pm 1\}$. Then, the normalized prolate uniaxial trace-free tensor in the direction of Ω , A_Ω , is defined by

$$A_\Omega = \Omega \otimes \Omega - \frac{1}{n} \text{Id}. \quad (44)$$

A_Ω is a traceless symmetric tensor with leading eigenvalue equal to $\frac{n-1}{n}$.

A_Ω is called a uniaxial tensor because it has only two eigenvalues with one being simple. The simple eigenvalue has associated normalized eigenvectors $\pm \Omega$. The line spanned by Ω is called the axis of the uniaxial tensor. It is trace-free and consequently, the two eigenvalues have opposite signs. It is called prolate because the simple eigenvalue is positive (it would be called oblate in the converse case). It is normalized meaning that its leading eigenvalue is exactly $\frac{n-1}{n}$. We note that A_Ω is invariant by the change $\Omega \rightarrow -\Omega$ showing that it actually depends on Ω seen as an element of the projective space $\mathbb{P}^{n-1} = \mathbb{S}^{n-1}/\{\pm 1\}$.

Proposition 1 (Gibbs distributions of uniaxial tensors). *The Gibbs distributions $G_{\eta A_\Omega}$ associated to tensors of the form ηA_Ω with $\eta > 0$ are given by*

$$G_{\eta A_\Omega}(\omega) = \frac{1}{Z_\eta} e^{\eta(\omega \cdot \Omega)^2}, \quad Z_\eta = \int_{\mathbb{S}^{n-1}} e^{\eta(\omega \cdot \Omega)^2} d\omega, \quad (45)$$

where the normalization constant Z_η does not depend on Ω but only on η .

Proof. Eq. (45) is obvious from (44). Defining $\theta \in (0, \pi)$ such that $\cos \theta = (\omega \cdot \Omega)$ and changing ω to (θ, z) where $z \in \mathbb{S}^{n-2}$ through $\omega = \cos \theta \Omega + \sin \theta z$, with $d\omega = C_n \sin^{n-2} \theta d\theta dz$ (C_n being such that $C_n \int_0^\pi \sin^{n-2} \theta d\theta = 1$ and $\int_{\mathbb{S}^{n-2}} dz = 1$), we get:

$$Z_\eta = C_n \int_0^\pi e^{\eta \cos^2 \theta} \sin^{n-2} \theta d\theta,$$

which does not depend on Ω . □

For two functions g and $\varphi: \mathbb{S}^{n-1} \rightarrow \mathbb{R}$, with $\varphi > 0$ a.e., we define:

$$\langle g \rangle_\varphi = \frac{\int_{\mathbb{S}^{n-1}} g(\omega) \varphi(\omega) d\omega}{\int_{\mathbb{S}^{n-1}} \varphi(\omega) d\omega}$$

We introduce the following

Definition 3.4 (Definition of S_2 and S_4). The quantities $S_2(\eta)$ and $S_4(\eta)$ are defined by

$$S_2(\eta) = \langle P_2(\omega \cdot \Omega) \rangle_{G_{\eta A_\Omega}}, \quad S_4(\eta) = \langle P_4(\omega \cdot \Omega) \rangle_{G_{\eta A_\Omega}}, \quad (46)$$

where $P_2(X)$ and $P_4(X)$ are the polynomials

$$\begin{aligned} P_2(X) &= \frac{1}{n-1} (nX^2 - 1), \\ P_4(X) &= \frac{1}{(n-1)(n+1)} [3 - 6(n+2)X^2 + (n+2)(n+4)X^4]. \end{aligned} \quad (47)$$

For the same reason as in Proposition 1, S_2 and S_4 do not depend on Ω . In dimension $n = 3$, the polynomials P_2 and P_4 are the Legendre polynomials of degree 2 and 4 respectively. About S_2 , we have the following proposition, which will be proved in Appendix B.1.

Proposition 2 (Properties of S_2). (i) We have

$$Q_{G_{\eta A_\Omega}} = S_2(\eta) A_\Omega. \quad (48)$$

(ii) The order parameter (12) of the distribution $\rho G_{\eta A_\Omega}$ is $\chi_{\rho G_{\eta A_\Omega}} = S_2(\eta)$.

(iii) S_2 is a non-decreasing function from $(0, \infty)$ onto $(0, 1)$, i.e. $S_2(0) = 0$ and $S_2 \rightarrow 1$ as $\eta \rightarrow \infty$.

We note that, when $\eta \rightarrow 0$, $G_{\eta A_\Omega}$ converges to the uniform probability distribution on \mathbb{S}^{n-1} . Likewise, when $\eta \rightarrow \infty$, $G_{\eta A_\Omega}$ concentrates on two Dirac deltas $\frac{1}{2}(\delta_\Omega + \delta_{-\Omega})$ which characterizes fully aligned distributions of molecules in the direction Ω . Therefore, S_2 takes the value 0 on fully disordered distributions and the value 1 on fully ordered ones. As η increases, $G_{\eta A_\Omega}$ shows increasing order evidenced by the increase of the order parameter S_2 . Now, we have the following

Proposition 3 (Implicit definition of $\eta(\rho)$). The implicit equation

$$\frac{\eta}{\alpha \rho} = S_2(\eta), \quad (49)$$

has at least a root η if and only if $\rho \in (\rho^*, +\infty)$ where $\rho^* > 0$. It has at most two roots. By choosing the largest root (which is necessarily nonnegative), it defines a smooth non-decreasing function $(\rho^*, +\infty) \rightarrow (\eta^*, +\infty)$, $\rho \mapsto \eta(\rho)$, where $\eta^* = \lim_{\rho \rightarrow \rho^*} \eta(\rho) \geq 0$.

This proposition is a consequence of the result of Wang and Hoffman [56] which will be recalled in Section 4. With this, we formulate the following conjecture, which has been verified in dimension $n = 2$ [28], $n = 3$ [1, 29, 47, 48] and $n = 4$ [31].

Conjecture 1 (Stable anisotropic equilibria). The set \mathcal{E} of stable anisotropic equilibria (in the sense of Remark 1) is given by

$$\mathcal{E} = \{\rho G_{\eta(\rho) A_\Omega} \mid \rho \in (\rho^*, +\infty), \Omega \in \mathbb{P}^{n-1}\}.$$

We will only consider anisotropic equilibria, i.e. belonging to the set \mathcal{E} above. Stable isotropic equilibria (i.e. such that $f = \rho$ is independent of ω) do exist but will not be used here.

Remark 2. In the case $n = 3$, using the change of variables $z = \cos \theta$ and an integration by parts, Eq. (49) can be recast as

$$\frac{3e^\eta}{\int_0^1 e^{\eta z^2} dz} = 3 + 2\eta + \frac{4\eta^2}{\alpha \rho}.$$

Upon changing η into $-\eta$ and making $\rho = 1$, we recover Eq. (1.9) of [48] and Eq. (3.2) of [59] (up to a typo in the latter: a factor 4 is missing in front of the η^2 term).

Now, we introduce the molecular interaction potential at equilibrium $U_{\rho G_{\eta A_\Omega}}^0$ where U_f^0 is given by (29). Thanks to (48), (49), we have

$$\alpha \rho Q_{G_{\eta A_\Omega}} = \eta(\rho) A_\Omega. \quad (50)$$

Thus, introducing $\theta \in [0, \pi]$ such that $\omega \cdot \Omega = \cos \theta$, straightforward computations give

$$\begin{aligned} U_{\rho G_{\eta A_{\Omega}}}^0 &= -\eta(\omega \cdot A_{\Omega}\omega) + \frac{n-1}{n}\alpha\rho = -\eta\left((\omega \cdot \Omega)^2 - \frac{1}{n}\right) + \frac{n-1}{n}\alpha\rho \quad (51) \\ &= -\eta\left(\cos^2 \theta - \frac{1}{n}\right) + \frac{n-1}{n}\alpha\rho =: \tilde{U}_0(\theta), \end{aligned}$$

so defining the function $\tilde{U}_0(\theta)$. We note that

$$\frac{d\tilde{U}_0}{d\theta} = 2\eta \cos \theta \sin \theta. \quad (52)$$

For two functions φ and ψ defined on $[0, \pi]$ with $\psi > 0$, a.e., we define

$$\langle\langle \varphi \rangle\rangle_{\psi} = \frac{\int_0^{\pi} \varphi(\theta) \psi(\theta) \sin^{n-2} \theta \, d\theta}{\int_0^{\pi} \psi(\theta) \sin^{n-2} \theta \, d\theta}.$$

Thanks to these notations, we can state the

Definition 3.5 (Auxiliary function g). The function $g: [0, \pi] \rightarrow \mathbb{R}$, $\theta \mapsto g(\theta)$, is the unique solution (in a sense made precise in Section 5) of the elliptic equation

$$\frac{1}{\sin^{n-2} \theta} \frac{d}{d\theta} \left(\sin^{n-2} \theta \frac{dg}{d\theta} \right) - \frac{d\tilde{U}_0}{d\theta} \frac{dg}{d\theta} - (n-2) \frac{g}{\sin^2 \theta} = -\frac{d\tilde{U}_0}{d\theta}. \quad (53)$$

Note that, in the special case $n = 3$, (53) coincides with Eq. (5.31) of [37]. Thanks to g we have the following proposition, proved in Section 6.2:

Proposition 4 (Constant c). Assume $\Lambda \neq 0$. Then, the constant c given by

$$c = \frac{(n-1)\Lambda S_2(\eta)}{\langle\langle g \frac{d\tilde{U}_0}{d\theta} \rangle\rangle_{\exp(\eta \cos^2 \theta)}}, \quad (54)$$

is such that $c/\Lambda > 0$.

In dimension $n = 3$, this formula coincides with formula (5.33) of [37]. We now introduce the following definitions

Definition 3.6 (Definition of the Leslie constants α_k , $k = 1, \dots, 6$). The Leslie constants α_k , $k = 1, \dots, 6$ are defined by

$$\alpha_1 = (\zeta - \Lambda^2)S_4, \quad \alpha_2 = -\frac{\Lambda S_2}{2} \left(\frac{1}{c} + 1 \right), \quad \alpha_3 = \frac{\Lambda S_2}{2} \left(\frac{1}{c} - 1 \right), \quad (55)$$

$$\alpha_4 = \frac{2(\zeta - \Lambda^2)}{(n+2)(n+4)} S_4 - \frac{2}{n} \left(\frac{\Lambda^2}{2} + \frac{2(\zeta - \Lambda^2)}{n+4} \right) S_2 + \frac{1}{n} \left(\Lambda^2 + \frac{2(\zeta - \Lambda^2)}{n+2} \right), \quad (56)$$

$$\alpha_5 = -\frac{2(\zeta - \Lambda^2)}{n+4} S_4 + \left(\frac{\Lambda}{2} + \frac{\Lambda^2}{2} + \frac{2(\zeta - \Lambda^2)}{n+4} \right) S_2, \quad (57)$$

$$\alpha_6 = -\frac{2(\zeta - \Lambda^2)}{n+4} S_4 + \left(-\frac{\Lambda}{2} + \frac{\Lambda^2}{2} + \frac{2(\zeta - \Lambda^2)}{n+4} \right) S_2, \quad (58)$$

where S_2 and S_4 are given by (46) and their dependence on η has been omitted for simplicity, and where c is given by (54). We note Parodi's relation: $\alpha_6 - \alpha_5 = \alpha_2 + \alpha_3$.

3.2. Main result: Statement and comments. Now, our aim is to prove the following formal result:

Theorem 3.7 (Formal limit of model (34) - (36)). *We assume $n \geq 2$, $\Lambda \neq 0$. For $n \geq 5$, we assume that Conjecture 1 is true (for $2 \leq n \leq 4$, this conjecture is a theorem [1, 28, 29, 31, 47, 48]). When $\varepsilon \rightarrow 0$, we assume that $(f^\varepsilon, u^\varepsilon) \rightarrow (f, u)$ as smoothly as needed, where $f(x, \cdot, t)$ is a stable anisotropic local equilibrium for all (x, t) . Then, on the open set*

$$\mathcal{B} = \{(x, t) \in \mathbb{R}^n \times [0, \infty) \mid \rho_f(x, t) > \rho^*\}, \quad (59)$$

(where ρ^* is defined at Proposition 3), we have

$$f(x, \omega, t) = \rho(x, t)G_{\eta(\rho(x, t))A_{\Omega(x, t)}}(\omega), \quad (60)$$

where the function $(\rho^*, \infty) \ni \rho \mapsto \eta(\rho) \in [0, \infty)$ is defined by (49). The functions $(x, t) \mapsto (\rho, \Omega, u)(x, t)$ satisfy the following system of partial differential equations (called the Ericksen-Leslie system):

$$\partial_t \rho + \nabla_x \cdot (\rho u) = 0, \quad (61)$$

$$\partial_t \Omega + u \cdot \nabla_x \Omega + W\Omega - c P_{\Omega^\perp} \left(E\Omega + \frac{2\beta}{\Lambda} \Delta_x(\eta\Omega) \right) = 0, \quad (62)$$

$$\partial_t u + u \cdot \nabla_x u + \nabla p = \frac{1}{\text{Re}} \left(\Delta_x u + \frac{1}{\text{Er}} \nabla_x \cdot \sigma \right), \quad (63)$$

$$\nabla_x \cdot u = 0, \quad (64)$$

$$\sigma = \sigma_L + \sigma_E, \quad (65)$$

$$\begin{aligned} \sigma_L = \rho \{ & \alpha_1 (E : (\Omega \otimes \Omega)) \Omega \otimes \Omega + \alpha_2 \Omega \otimes N + \alpha_3 N \otimes \Omega \\ & + \alpha_4 E + \alpha_5 (\Omega \otimes \Omega) E + \alpha_6 E (\Omega \otimes \Omega) \}, \end{aligned} \quad (66)$$

$$\begin{aligned} \sigma_E = & -\frac{2\beta}{\alpha} \nabla_x(\eta\Omega) \nabla_x(\eta\Omega)^T \\ & + \frac{(n+1)\beta}{n\alpha} \nabla_x \eta \otimes \nabla_x \eta + \frac{(n-1)\alpha\beta}{n} \nabla_x \rho \otimes \nabla_x \rho, \end{aligned} \quad (67)$$

where W and E are given by (3), β by (28), c by (54), α_k , $k = 1, \dots, 6$ by (55)-(58), and N by

$$N = D_t \Omega + W\Omega, \quad (68)$$

with D_t given by (23).

Remark 3. Using (49), we have the following equivalent expression of σ_E :

$$\sigma_E = -\frac{2\beta}{\alpha} \nabla_x \Omega \nabla_x \Omega^T - \frac{(n-1)\beta}{n\alpha} \left[1 - \frac{1}{S_2^2} \left(1 - \eta \frac{S_2'}{S_2} \right)^2 \right] \nabla_x \eta \otimes \nabla_x \eta,$$

where S_2' denotes the derivative of S_2 with respect to η . In particular, this formula shows that the contribution of the density gradient to σ_E is a rank-1 tensor (which is not obvious from (67); on the other hand, (67) has more symmetry between $\nabla_x \rho$ and $\nabla_x \eta$).

Remark 4. In the case $\Lambda = 0$, the result is still valid, except that (62) must be replaced by

$$\partial_t \Omega + u \cdot \nabla_x \Omega + W\Omega - 2\beta \tilde{c} P_{\Omega^\perp} \Delta_x(\eta\Omega) = 0,$$

where $\tilde{c} = (n-1)S_2(\eta) / \langle g \frac{d\tilde{U}_0}{d\theta} \rangle_{\exp(\eta \cos^2 \theta)}$.

In the literature [37, 59], Eq. (62) is written differently. For this we need the

Definition 3.8 (Molecular field and γ -constants). We define

$$\begin{aligned}\gamma_1 &= \frac{\Lambda S_2}{c} = \alpha_3 - \alpha_2, & \gamma_2 &= -\Lambda S_2 = \alpha_6 - \alpha_5 = \alpha_2 + \alpha_3, \\ H &= 2\beta S_2 \Delta_x(\eta\Omega).\end{aligned}\quad (69)$$

The quantity H is called the molecular field.

Then, we have the following proposition, whose proof is immediate:

Proposition 5 (Equivalent form of Eq. (62)). *Eq. (62) is equivalent to*

$$P_{\Omega^\perp}(H - \gamma_1 N - \gamma_2 E\Omega) = 0. \quad (70)$$

We compare System (61)-(67) with the literature. Ref. [37] considers a spatially homogeneous model in dimension $n = 3$ with $\zeta = 0$. Spatial homogeneity means that ρ and Ω do not depend on x , and so $H = 0$, $\sigma_E = 0$ and $N = \partial_t \Omega + W\Omega$ while E and W are constant. In this case, our model reduces to (70) (with $H = 0$) and $\sigma = \sigma_L$ with σ_L given by (66), which are the two equations obtained in [37], provided the external magnetic field considered in [37] is set to 0. Finally, formulas (55)-(58) for $n = 3$ and $\zeta = 0$ are identical with Formula (6.2) of [37]. So, our model is consistent with [37].

Then, Refs. [25, 59] consider a spatially non-homogeneous setting, but still with a constant and uniform ρ (we easily see that $\rho = \text{Constant}$ is consistent with both the kinetic model (34) and the fluid one (61) due to the incompressibility conditions (36) and (64)). Their setting is $n = 3$, $\zeta = \frac{1}{2}$ and $\Lambda = 1$. In this case, we see that formulas (55)-(58) are identical with Formulas (2.6), (2.7) of [59]. If $\rho = \text{Constant}$, then, $\eta = \text{Constant}$ as well. So, the Ericksen stresses and molecular field reduce to

$$\sigma_E = -k \nabla_x \Omega \nabla_x \Omega^T, \quad \rho H = k \Delta_x \Omega, \quad \text{with } k = \frac{2\beta}{\alpha} \eta^2, \quad (71)$$

which are the corresponding expressions (see top of p. 7) of [59]. With these expressions, our model reduces to (5) coupled with (63)-(66) and (71). It is identical with the model obtained in [59].

So, our model is consistent with the literature but has two additional features: the consideration of an arbitrary dimension $n \geq 2$ and the spatial non-homogeneity of ρ (and consequently, of η) which brings additional components to the elastic stresses and, as we will see below, to the elastic energy. Non-uniform η has been previously considered in [9, 8, 27, 40, 41, 46], but to the best of our knowledge, none has explicitly linked it to the polymer density and to kinetic theory.

A well-posedness theory of System (61)-(67) is outside the scope of this paper (see e.g. [35, 42, 43, 44, 60] for existence results of the Ericksen-Leslie system in a variety of forms). Note however that a condition for the well-posedness of the parabolic equation (62) is that $\frac{\beta c}{\Lambda} > 0$. This is indeed ensured by Prop. 4.

The main objective of this paper is to provide a (formal) derivation of Eqs. (61), (62) using the moment method and the generalized collision invariant (GCI) concept. Prior to this, in Section 4, we will return to the determination of the stable equilibria of the Doi model and provide support to Conjecture 1 and to Formula (49) linking ρ and η . Then, in Section 5, we develop the GCI concept and discuss its rationale and how it can be linked to the Hilbert expansion procedure. The derivation of (62) itself will be performed in Section 6. The second main objective of the paper is to provide expressions for the Leslie and Ericksen stresses in arbitrary dimension and for spatially inhomogeneous densities, which, to the

best of our knowledge, has not been considered before. As these computations are lengthy, they are deferred to Appendix B. Other auxiliary results can be found in this appendix and in the subsequent ones, Appendices C and D.

3.3. Energetics of the Ericksen-Leslie system. Next, we define the following energies:

Definition 3.9 (Oseen-Franck and Ericksen-Leslie energies). (i) The Oseen-Franck energy is defined by:

$$\begin{aligned}\mathcal{E}_F &= \frac{2\beta}{\alpha} \int_{\mathbb{R}^n} \frac{|\nabla_x(\eta\Omega)|^2}{2} dx - \alpha\beta \frac{n-1}{n} \int_{\mathbb{R}^n} \frac{|\nabla_x\rho|^2}{2} dx - \beta \frac{n+1}{n\alpha} \int_{\mathbb{R}^n} \frac{|\nabla_x\eta|^2}{2} dx \\ &=: \mathcal{E}_F^\Omega + \mathcal{E}_F^\rho + \mathcal{E}_F^\eta.\end{aligned}\tag{72}$$

(ii) The Ericksen-Leslie energy is defined by

$$\mathcal{E}_{EL} = \int_{\mathbb{R}^n} |u|^2 dt + \frac{1}{\text{ReEr}} \mathcal{E}_F.$$

Remark 5. (i) If ρ is uniformly constant (and hence, η too), \mathcal{E}_F reduces to

$$\mathcal{E}_F = \frac{2\beta\eta^2}{\alpha} \int_{\mathbb{R}^n} \frac{|\nabla_x\Omega|^2}{2} dx,$$

which is the classical Oseen-Franck elastic energy [25, 59]. The additional terms \mathcal{E}_F^ρ and \mathcal{E}_F^η make up for the non-uniformity of ρ and η .

(ii) Using (49), we find an alternate expression of \mathcal{E}_F :

$$\mathcal{E}_F = \frac{2\beta}{\alpha} \int_{\mathbb{R}^n} \eta^2 \frac{|\nabla_x\Omega|^2}{2} dx + \frac{(n-1)\beta}{n\alpha} \int_{\mathbb{R}^n} \left(1 - \frac{1}{S_2^2} (1 - \eta \frac{S_2'}{S_2})^2\right) \frac{|\nabla_x\eta|^2}{2} dx.$$

In particular, we see that this energy is positive if the following relation holds

$$1 - \frac{1}{S_2^2} (1 - \eta \frac{S_2'}{S_2})^2 \geq 0.$$

The investigation of this property is left to future work.

Now, we have the following proposition, which relates the molecular field to the derivative of the Franck energy with respect to the orientation field Ω .

Proposition 6 (Relation between the Franck energy and the molecular field). *We have the following relation:*

$$\rho H = -\frac{\delta\mathcal{E}_F}{\delta\Omega}(\eta, \Omega) = \frac{2\beta}{\alpha} \eta \Delta_x(\eta\Omega),\tag{73}$$

where $\frac{\delta\mathcal{E}_F}{\delta\Omega}(\eta, \Omega)$ is the functional derivative of \mathcal{E}_F with respect to the field Ω evaluated at the pair (η, Ω) .

Proof. For a $n \times n$ tensor S , we introduce the following energy density

$$e_F^\Omega(S) = \frac{2\beta}{\alpha} \frac{|S|^2}{2},$$

so that we can write

$$\mathcal{E}_F^\Omega = \int_{\mathbb{R}^n} e_F^\Omega(\nabla_x(\eta\Omega)) dx.$$

Now, straightforward computations show that the functional derivative $\frac{\delta \mathcal{E}_F}{\delta \Omega}$ is given by

$$\frac{\delta \mathcal{E}_F}{\delta \Omega}(\eta, \Omega) = \frac{\delta \mathcal{E}_F^\Omega}{\delta \Omega}(\eta, \Omega) = -\eta \nabla_x \cdot \left(\frac{\partial \mathcal{E}_F^\Omega}{\partial S}(\nabla_x(\eta \Omega)) \right) = -\frac{2\beta}{\alpha} \eta \Delta_x(\eta \Omega) = -\rho H,$$

where the first equality is due to the fact that the energies \mathcal{E}_F^ρ and \mathcal{E}_F^η do not depend on Ω , and the last one, to (49). Then, Eq. (73) follows. \square

The following proposition gives the energy identity for the Ericksen-Leslie system. Its proof is developed in Appendix B.4

Proposition 7 (Energy identity for the Ericksen-Leslie system). *We have the following identity:*

$$\begin{aligned} \frac{d\mathcal{E}_{EL}}{dt} + \mathcal{D}_{EL} &= 0, \\ \mathcal{D}_{EL} &= \frac{1}{\text{Re}} \int_{\mathbb{R}^n} |\nabla_x u|^2 dx + \frac{1}{\text{ReEr}} \int_{\mathbb{R}^n} \rho \left\{ \left(\alpha_1 + \frac{\gamma_2^2}{\gamma_1} \right) (E : (\Omega \otimes \Omega))^2 + \alpha_4 |E|^2 \right. \\ &\quad \left. + \left(\alpha_5 + \alpha_6 - \frac{\gamma_2^2}{\gamma_1} \right) |E\Omega|^2 + \frac{1}{\gamma_1} |P_{\Omega^\perp} H|^2 \right\} dx. \end{aligned} \quad (74)$$

Remark 6. (i) The use of this energy identity to derive a priori bounds for the solution of the Ericksen-Leslie equations is subject to two conditions: first, that the Oseen-Frank energy is positive as already mentioned in Remark 5; second, that the dissipation functional \mathcal{D}_{EL} is positive as well, which is not obvious given that the coefficients are not all positive. In [59], it is shown that, in the case $n = 3$, $\zeta = \frac{1}{2}$ and $\Lambda = 1$, \mathcal{D}_{EL} is positive. Besides, conditions for the positive-definiteness of \mathcal{D}_{EL} with coefficients which are not necessarily linked with a microscopic model can be found in [60]. The inspection of the positivity of \mathcal{E}_F and \mathcal{D}_{EL} for the present model is left to future work.

(ii) It is expected that this energy identity is the limit as $\varepsilon \rightarrow 0$ of the free-energy dissipation identity (24) of the Doi-Navier-Stokes system. This is indeed formally shown in [25]. However, due to the presence of the square of the Deborah number at the denominator of (27), we expect that the limiting free-energy dissipation identity will involve the first order correction $f^1 = \lim_{\varepsilon \rightarrow 0} \frac{f^\varepsilon - f^0}{\varepsilon}$. Showing that the terms involving f^1 eventually vanish is not obvious and left to future work.

4. Local equilibria. In this section, we develop the rationale for Conjecture 1. Since we aim at formal convergence results only, we suppose that the solution f^ε to (40) satisfies

$$f^\varepsilon \rightarrow f \quad \text{as } \varepsilon \rightarrow 0 \quad \text{as smoothly as needed.}$$

Then, from (40), it follows that f should satisfy (42), i.e. should be an equilibrium for any (x, t) . Eq. (42) leaves the dependence of f on (x, t) undetermined. Such an equilibrium is called ‘local’ (by contrast to a global equilibrium where f should not depend on (x, t)).

In this section, our goal is to determine the stable equilibria. Indeed, we anticipate that only stable equilibria can lead to a long time dynamics described by hydrodynamic equations. First, we should note that local equilibria are known in any dimension n [56] (see also [15, 28, 49] for the case $n = 2$ and [1, 14, 29, 47, 48, 61, 63] for the case $n = 3$). However, the stability of these equilibria is not known for general dimension n but only for $n = 2$ [28], $n = 3$ [1, 29, 47, 48] and $n = 4$ [31].

These results strongly support a conjecture about the stable equilibria in general dimension n that we will make below and whose rigorous investigation is deferred to future work. We first need to introduce a set of notations and intermediate results.

Definition 4.1 (Auxiliary operator). Let S be a trace-free symmetric matrix. Then, the auxiliary operator L_S is given by

$$L_S f = \nabla_\omega \cdot \left[G_S \nabla_\omega \left(\frac{f}{G_S} \right) \right]. \tag{75}$$

with G_S given by (43).

The relation between the collision operator $C(f)$ and the auxiliary operator L_S is given by the following lemma. Note that L_S is NOT the linearization of C about G_S .

Lemma 4.2 (Relation between C and L). *We have*

$$C(f) = L_{\alpha \rho_f Q_f} f. \tag{76}$$

Proof of Lemma 4.2. We can write

$$L_S f = \nabla_\omega \cdot \left[\nabla_\omega f - f \nabla_\omega (\log G_S) \right].$$

But $-\log G_S = -\omega \cdot S\omega + \log Z_S$. So, $-\log G_{\alpha \rho_f Q_f} = U_f^0 + \tilde{Z}(f)$ where $\tilde{Z}(f)$ does not depend on ω . Thus, $-\nabla_\omega (\log G_{\alpha \rho_f Q_f}) = \nabla_\omega U_f^0$ and so, $L_{\alpha \rho_f Q_f} = C(f)$, thanks to (39). \square

Now, we have a first result:

Lemma 4.3 (First step towards a characterization of the equilibria). *(i) Let $f \geq 0$, $f \neq 0$ be an equilibrium. Then, there exists $\rho > 0$ and a trace-free symmetric matrix Q such that*

$$f = \rho G_{\alpha \rho Q}. \tag{77}$$

(ii) Reciprocally, let f be given by (77). Then, f is an equilibrium if and only if Q satisfies the fixed-point equation also known as the compatibility equation:

$$Q = Q_{\rho G_{\alpha \rho Q}}, \tag{78}$$

where we recall that for a distribution f , Q_f is given by (10).

Proof. (i) Suppose $C(f) = 0$. Letting $S = \alpha \rho_f Q_f$, (76) implies $L_S f = 0$. Multiplying (75) by f/G_S , integrating over \mathbb{S}^{n-1} and using Green's formula leads to

$$\int_{\mathbb{S}^{n-1}} G_S \left| \nabla_\omega \left(\frac{f}{G_S} \right) \right|^2 d\omega = 0.$$

Since the quantity inside the integral is nonnegative, and $G_S > 0$, this implies $\nabla_\omega \left(\frac{f}{G_S} \right) = 0$. So, there exists $\rho > 0$ such that $f = \rho G_S$ which leads to (77).

(ii) Let f be given by (77). Then, since $G_{\alpha \rho Q}$ is a probability density, we have $\rho_f = \rho$. Now, from the proof of Part (i), if f is an equilibrium, then $f = \rho_f G_{\alpha \rho_f Q_f}$. We deduce that $G_{\alpha \rho Q_f} = G_{\alpha \rho Q}$, and, by taking the logarithm, that

$$\omega \cdot (Q_f - Q)\omega = \frac{1}{\alpha \rho} (\log Z_{\alpha \rho Q_f} - \log Z_{\alpha \rho Q}) =: \mu,$$

where μ is a constant, independent of ω . So, $Q_f - Q - \mu \text{Id}$ is the matrix of a quadratic form which is zero on \mathbb{S}^{n-1} and so, by homogeneity, on \mathbb{R}^n . Thus, $Q_f - Q - \mu \text{Id} = 0$

and, owing to the fact that Q_f and Q are trace-free, we have $\mu = 0$. It follows that $Q_f = Q$. Replacing f by its expression (77), we get (78). \square

To complete the characterization of the equilibria, we need to solve the compatibility equation (78). As pointed out above, this has been done in any dimension n in [56] (see also [49] for $n = 2$ and [1, 29, 47, 48, 63] for $n = 3$). This result is summarized without proof in the following lemma

Lemma 4.4 (Final characterization of the equilibria [56]). *Let f be an equilibrium. Then Q_f has at most two distinct eigenvalues.*

- *If all eigenvalues of Q_f are identical, then $Q_f = 0$ and $f = \rho$ is a uniform equilibrium.*
- *If Q_f has exactly two distinct eigenvalues, denote by λ_f its largest eigenvalue and by \mathcal{Y}_f the associated eigenspace, supposed of dimension d such that $1 \leq d \leq n - 1$. Then, $0 < \lambda_f < \frac{1}{d} - \frac{1}{n}$ and Q_f is written*

$$Q_f = B_{\lambda_f, \mathcal{Y}_f} := \lambda_f \left(P_{\mathcal{Y}_f} - \frac{d}{n-d} P_{\mathcal{Y}_f^\perp} \right), \tag{79}$$

where $P_{\mathcal{Y}_f}$ and $P_{\mathcal{Y}_f^\perp}$ are the orthogonal projections of \mathbb{R}^n onto \mathcal{Y}_f and \mathcal{Y}_f^\perp respectively. Then, f is of the form

$$f = \rho_d^n(\lambda_f) G_{\alpha \rho_d^n(\lambda_f) B_{\lambda_f, \mathcal{Y}_f}},$$

where $\rho_d^n: [0, \frac{1}{d} - \frac{1}{n}] \rightarrow [0, \infty)$, $\lambda \mapsto \rho_d^n(\lambda)$ is a specific function (not detailed here except for the case $d = 1$, see below). Furthermore, λ_f is a root of the equation

$$\rho_d^n(\lambda) = \rho. \tag{80}$$

The existence and number of classes of equilibria such that $\rho_f = \rho$ are determined by the existence and number of roots λ of Eq. (80). A given root λ gives rise to a family of equilibria parametrized by the Grassmann manifold $Gr(k, n)$ of d -dimensional vector subspaces \mathcal{Y} of \mathbb{R}^n .

Here, we are only interested in the case $d = 1$ as we will conjecture that this is the only case which includes stable equilibria (see conjecture 2 below). For simplicity, ρ^n stands for the function ρ_1^n . In the case $n = 2$, ρ^2 is monotonously increasing and maps $[0, \frac{1}{2})$ onto the interval $[\rho^*, +\infty)$ with $\rho^* = \rho^2(0)$ (see Fig. 1a). In the case $n \geq 3$, ρ^n is decreasing in the interval $[0, \lambda^*]$ and increasing in $[\lambda^*, 1 - \frac{1}{n})$. Thus $\rho^* = \rho(\lambda^*)$ is a global minimum of ρ^n (see Fig. 1b). In all cases, the equation $\rho_n(\lambda) = \rho$ has a solution if and only if $\rho \geq \rho^*$ and this solution is unique in the case $n = 2$ while, in the case $n \geq 3$, there are two solutions if $\rho \in (\rho^*, \rho^n(0)]$, and one solution if $\rho \in \{\rho^*\} \cup (\rho^n(0), \infty)$ (see [56] for details).

As already stated, for general n , the stability of the equilibria described in Lemma 4.4 is not known yet. However, their stability is known for $n = 2$ [28], $n = 3$ [1, 29, 47, 48] and $n = 4$ [31]. Based on these results, we formulate the following conjecture for any dimension $n \geq 2$ and refer to the above-mentioned references for details on the notion of stability involved.

Conjecture 2 (Stable anisotropic equilibria). *For any dimension $n \geq 2$, the branch of solutions to the equation $\rho^n(\lambda) = \rho$ (which corresponds to $d = 1$) with largest λ , which is defined for $\rho \in (\rho^*, \infty)$, corresponds to the unique class of stable anisotropic equilibria.*




FIGURE 1. Graphical representation of the function $\lambda \mapsto \rho^n(\lambda)$ (after [56]). (a) case $n = 2$. (b) case $n \geq 3$. The portions of the curves that correspond to stable equilibria are in blue, the unstable ones, in green.

We denote by the function $\lambda: (\rho^*, +\infty) \rightarrow (\lambda^*, 1 - \frac{1}{n})$, $\rho \mapsto \lambda(\rho)$, the largest solution to $\rho^n(\lambda) = \rho$. With Conjecture 2, the stable equilibria f correspond to the class of equilibria described in Lemma 4.4, Case 2, with $d = 1$ and $\lambda_f = \lambda(\rho)$. In this case, \mathcal{Y}_f is one-dimensional and thus, spanned by a unique normalized vector (up to a sign) $\Omega \in \mathbb{P}^{n-1}$. Hence, we have $P_{\mathcal{Y}_f} = \Omega \otimes \Omega$ and $P_{\mathcal{Y}_f^\perp} = P_{\Omega^\perp}$. Then by (79),

$$Q_f = \frac{n}{n-1} \lambda(\rho) A_\Omega, \tag{81}$$

where A_Ω is the normalized uniaxial tensor given by (44). Defining

$$\eta(\rho) = \frac{n}{n-1} \alpha \rho \lambda(\rho), \tag{82}$$

from (78) we get that the equilibria are of the form $f = \rho G_{\eta(\rho)A_\Omega}$ where ρ is arbitrary as long as $\lambda(\rho)$ is defined, i.e. $\rho \in (\rho^*, \infty)$, and where Ω is arbitrary in \mathbb{P}^{n-1} . Hence, Conjecture 1 is a direct consequence of Conjecture 2, provided we show that the function $\rho \mapsto \eta(\rho)$ is the one given by Proposition 3, which we do now:

Proof of Proposition 3. Equating (81) with (48) and using (82), we get (49). The root with the largest η must be chosen because this corresponds to the choice of largest λ in Conjecture 2 (as λ is proportional to η by (82)). \square

From (40) and Conjecture 2, we deduce the:

Corollary 1 (Local equilibria). *Let f be the formal limit of f^ε as $\varepsilon \rightarrow 0$ and suppose that $u^\varepsilon \rightarrow u$ smoothly. On the open set \mathcal{B} defined by (59), f is given by (60) where $\rho = \rho_f: (x, t) \in \mathbb{R}^n \times [0, \infty) \mapsto [\rho^*, \infty)$ and $\Omega: (x, t) \in \mathbb{R}^n \times [0, \infty) \mapsto \mathbb{P}^{n-1}$ are functions such that f satisfies*

$$T_u(f) = \lim_{\varepsilon \rightarrow 0} \frac{C(f^\varepsilon)}{\varepsilon}. \tag{83}$$

Note that $\rho = \rho_f$ is the local density associated to f , while $\Omega(x, t)$ if the axis of the uniaxial Q-tensor Q_f thanks to (48). The restriction to the set \mathcal{B} is needed to ensure that $\eta(\rho(x, t))$ is well-defined. The determination of the functions (ρ, Ω) such that (83) holds is quite challenging, due to the presence of ε in the denominator at the right-hand side. It will require the Generalized Collision Invariant concept as detailed below.

5. Generalized collision invariants.

5.1. Collision invariant. We first recall the notion of Collision Invariant (CI). The goal is to eliminate the singular right hand side of (83) by using integration against appropriate test functions. More precisely we have:

Definition 5.1. A Collision Invariant (CI) $\psi(\omega)$ is a function such that

$$\int_{\mathbb{S}^{n-1}} C(f) \psi \, d\omega = 0, \quad \forall f.$$

Here, we do not specify any regularity requirement on ψ since our goal is to develop a formal theory only. If ψ is a CI, using it as a test function for (40), we have, after integration with respect to ω and omitting ε as the identity is valid for any ε :

$$\begin{aligned} \partial_t \left(\int_{\mathbb{S}^{n-1}} f \psi \, d\omega \right) + \nabla_x \cdot \left(u \int_{\mathbb{S}^{n-1}} f \psi \, d\omega \right) - \int_{\mathbb{S}^{n-1}} \nabla_\omega \psi \cdot (\Delta P_{\omega^\perp} E - W) \omega f \, d\omega \\ - 2\alpha \beta \int_{\mathbb{S}^{n-1}} \nabla_\omega \psi \cdot P_{\omega^\perp} \Delta_x (\rho_f Q_f) \omega f \, d\omega = 0, \end{aligned} \tag{84}$$

which is an evolution equation for the moment $\int_{\mathbb{S}^{n-1}} f \psi \, d\omega$. Since this equation does not depend on ε , it is still verified by the solution of (83). We have an obvious CI, namely, $\psi = 1$, which leads to the mass conservation (or continuity) equation

$$\partial_t \rho_f + \nabla_x \cdot (\rho_f u) = 0. \tag{85}$$

In particular, taking the limit $\varepsilon \rightarrow 0$, it shows (61). As u is divergence free thanks to (36), (85) can be equivalently written

$$D_t \rho_f = 0, \tag{86}$$

with D_t given by (23).

Any odd function ψ of ω is also a CI. However, it is not invariant when ω is changed into $-\omega$, a condition that has been enforced throughout this work (see e.g. (1)). Indeed, Eq. (84) with odd functions ψ have all their terms identically zero and do not provide any useful information. We do not have any other obvious CI. Therefore, we are lacking an equation for Ω . In order to overcome this problem, we use the concept of ‘‘Generalized Collision Invariant (GCI)’’ introduced in [23] and adapted to the present context.

5.2. Generalized collision invariant: Definition and characterization. To introduce the GCI concept, we first need some additional notations and definitions.

Definition 5.2 (and notations). (i) \mathcal{S}_n^0 is the vector space of symmetric trace free $n \times n$ matrices.

(ii) \mathcal{U}_n^0 is the subset of \mathcal{S}_n^0 consisting of tensors whose leading eigenvalue is equal to

$\frac{n-1}{n}$ and is simple.

(iii) We denote by λ_f the leading eigenvalue of Q_f and by η_f the following quantity:

$$\eta_f = \alpha \rho_f \frac{n}{n-1} \lambda_f. \quad (87)$$

From (8), we have

$$0 \leq \frac{n}{n-1} \lambda_f \leq 1.$$

Note that in general, λ_f may not be simple.

(iv) If $Q_f \neq 0$, then $\lambda_f \neq 0$ and we define the ‘‘Normalized Q-Tensor (NQT) of f ’’, Σ_f by

$$\Sigma_f = \frac{n-1}{n} \frac{Q_f}{\lambda_f}. \quad (88)$$

$\Sigma_f \in \mathcal{S}_n^0$. Its leading eigenvalue is $\frac{n-1}{n}$ which, again, may not be simple.

(vi) Let $\Sigma \in \mathcal{U}_n^0$. We denote by $\Omega_\Sigma \in \mathbb{P}^{n-1}$ the normalized eigenvector (up to a sign) associated with the simple eigenvalue $\frac{n-1}{n}$ of Σ . Note that the tensor A_{Ω_Σ} is uniquely defined, irrespective of the choice of the sign of Ω_Σ .

(v) Suppose $\Sigma_f \in \mathcal{U}_n^0$. Then, Ω_{Σ_f} is simply denoted by Ω_f .

Remark 7. From (48), we get that $\Sigma_{\rho G_{\eta A_\Omega}} = A_\Omega$ meaning that the NQT’s of the stable anisotropic equilibria are all equal to A_Ω .

We recall that the auxiliary operator L_S for $S \in \mathcal{S}_n^0$ is defined by (75). The GCI are now defined in the following

Definition 5.3. Let $(\eta, \Sigma) \in (0, \infty) \times \mathcal{U}_n^0$. A Generalized Collisional Invariant (GCI) associated to the pair (η, Σ) is a function ψ such that

$$\int_{\mathbb{S}^{n-1}} (L_{\eta\Sigma} f) \psi \, d\omega = 0 \quad \text{for all } f \text{ such that } P_{\Omega_\Sigma^\perp}(Q_f \Omega_\Sigma) = 0. \quad (89)$$

The set of GCI associated to a given pair $(\eta, \Sigma) \in (0, \infty) \times \mathcal{U}_n^0$ is a linear vector space and is denoted by $\mathcal{C}_{\eta\Sigma}$.

There is a rationale for this definition, which is developed in Section 5.3 below.

The following lemma gives the equation satisfied by the GCI:

Lemma 5.4. Let $(\eta, \Sigma) \in (0, \infty) \times \mathcal{U}_n^0$. Then $\psi \in \mathcal{C}_{\eta\Sigma}$ if and only if there exists $V \in \{\Omega_\Sigma\}^\perp$ such that

$$\nabla_\omega \cdot (G_{\eta\Sigma}(\omega) \nabla_\omega \psi) = (\omega \cdot \Omega_\Sigma) (\omega \cdot V) G_{\eta\Sigma}(\omega), \quad \forall \omega \in \mathbb{S}^{n-1}. \quad (90)$$

Proof. For $\Omega \in \mathbb{S}^{n-1}/\{\pm 1\}$, we define the following space of functions:

$$\mathcal{X}_\Omega = \{\mathbb{S}^{n-1} \ni \omega \mapsto (\Omega \cdot \omega) (V \cdot \omega) \in \mathbb{R} \mid V \in \{\Omega\}^\perp\}, \quad (91)$$

The space \mathcal{X}_Ω is a finite-dimensional subspace of $L^2(\mathbb{S}^{n-1})$. We first note that for any $f \in L^2(\mathbb{S}^{n-1})$, we have

$$\begin{aligned} P_{\Omega^\perp}(Q_f \Omega) = 0 &\iff \int_{\mathbb{S}^{n-1}} f(\omega) (\omega \cdot V) (\omega \cdot \Omega) \, d\omega = 0, \quad \forall V \in \{\Omega\}^\perp \\ &\iff f \in \mathcal{X}_\Omega^\perp, \end{aligned} \quad (92)$$

where the orthogonality is meant with respect to the standard L^2 -product on $L^2(\mathbb{S}^{n-1})$.

On the other hand, we note that $\int_{\mathbb{S}^{n-1}} (L_{\eta\Sigma} f) \psi \, d\omega = 0$ is equivalent to saying that $f \in \{L_{\eta\Sigma}^* \psi\}^\perp$ where again, the orthogonality is meant with respect to the

standard L^2 -product on $L^2(\mathbb{S}^{n-1})$ and where $L_{\eta\Sigma}^*$ is the formal L^2 -adjoint of $L_{\eta\Sigma}$, i.e.

$$L_{\eta\Sigma}^*\psi = \frac{1}{G_{\eta\Sigma}} \nabla_\omega \cdot (G_{\eta\Sigma} \nabla_\omega \psi).$$

Therefore, thanks to (92), Condition (89) is equivalent to saying that

$$f \in \mathcal{X}_{\Omega_\Sigma}^\perp \implies f \in \{L_{\eta\Sigma}^*\psi\}^\perp,$$

or in other words, that $\mathcal{X}_{\Omega_\Sigma}^\perp \subset \{L_{\eta\Sigma}^*\psi\}^\perp$. Taking the orthogonal to this relation and noting that both $\mathcal{X}_{\Omega_\Sigma}$ and $\text{Span}\{L_{\eta\Sigma}^*\psi\}$ (where for a subset B of a vector space, $\text{Span } B$ denotes the subspace generated by B) are finite-dimensional, hence, closed subspaces of $L^2(\mathbb{S}^{n-1})$, we get $\text{Span}\{L_{\eta\Sigma}^*\psi\} \subset \mathcal{X}_{\Omega_\Sigma}$. In particular, this implies that there exists $V \in \{\Omega_\Sigma\}^\perp$ such that $L_{\eta\Sigma}^*\psi(\omega) = (\omega \cdot \Omega_\Sigma)(\omega \cdot V)$, which, upon multiplying by $G_{\eta\Sigma}$, gives (90). The converse is straightforward. \square

Now, we give an existence theory for the solutions of (90). We denote by $H^1(\mathbb{S}^{n-1})$ the space of square integrable functions of \mathbb{S}^{n-1} into \mathbb{R} whose derivatives are square integrable and introduce

$$\dot{H}^1(\mathbb{S}^{n-1}) = \left\{ u \in H^1(\mathbb{S}^{n-1}) \mid \int_{\mathbb{S}^{n-1}} u(\omega) d\omega = 0 \right\}.$$

Then we have the

Proposition 8. *Let $(\eta, \Sigma) \in (0, \infty) \times \mathcal{U}_n^0$ and $V \in \{\Omega_\Sigma\}^\perp$. Then, there exists a unique solution of (90) in $\dot{H}^1(\mathbb{S}^{n-1})$ denoted by $\psi_{\eta\Sigma, V}$. The linear vector space $\mathcal{C}_{\eta\Sigma}$ of GCI associated with (η, Σ) is given by*

$$\mathcal{C}_{\eta\Sigma} = \{C_0 + \psi_{\eta\Sigma, V} \mid C_0 \in \mathbb{R}, V \in \{\Omega_\Sigma\}^\perp\}. \tag{93}$$

Proof. We look for solutions of (90) in variational form. The variational formulation reads as follows: find $\psi \in H^1(\mathbb{S}^{n-1})$ such that

$$\int_{\mathbb{S}^{n-1}} G_{\eta\Sigma} \nabla_\omega \psi \cdot \nabla_\omega \theta d\omega = - \int_{\mathbb{S}^{n-1}} G_{\eta\Sigma} (\omega \cdot \Omega_\Sigma) (\omega \cdot V) \theta d\omega, \quad \forall \theta \in H^1(\mathbb{S}^{n-1}). \tag{94}$$

By Poincaré inequality and the fact that $G_{\eta\Sigma}$ is smooth and bounded from above and below, the bilinear form $\int G_{\eta\Sigma} \nabla_\omega \psi \cdot \nabla_\omega \theta d\omega$ is continuous and coercive on $\dot{H}^1(\mathbb{S}^{n-1})$. Therefore, by Lax-Milgram theorem, the variational formulation (94) has a unique solution in $\dot{H}^1(\mathbb{S}^{n-1})$ denoted by $\psi_{\eta\Sigma, V}$ when θ is restricted to belong to $\dot{H}^1(\mathbb{S}^{n-1})$. To show that this is a solution for all $\theta \in H^1(\mathbb{S}^{n-1})$, it is enough to show that it satisfies (94) for $\theta = 1$, i.e. that the following holds:

$$\int_{\mathbb{S}^{n-1}} G_{\eta\Sigma} (\omega \cdot \Omega_\Sigma) (\omega \cdot V) d\omega = 0, \quad \forall V \in \{\Omega_\Sigma\}^\perp. \tag{95}$$

Let (e_1, \dots, e_n) with $e_n = \Omega_\Sigma$ be an ortho-normal basis of \mathbb{R}^n consisting of eigenvectors of Σ . Let $\lambda_1, \dots, \lambda_n$ be the associated eigenvalues. Let $\omega = \sum_{k=1}^n \omega_k e_k$ be the decomposition of ω in this basis. It is enough to show (95) for $V = e_j$ with $j \in \{1, \dots, n-1\}$. Then, we have

$$\int_{\mathbb{S}^{n-1}} G_{\eta\Sigma} (\omega \cdot \Omega_\Sigma) (\omega \cdot e_j) d\omega = \frac{1}{Z_{\eta\Sigma}} \int_{\mathbb{S}^{n-1}} e^{\eta(\lambda_1 \omega_1^2 + \dots + \lambda_n \omega_n^2)} \omega_j \omega_n d\omega = 0,$$

thanks to the change of ω_n into $-\omega_n$. This shows (95) and so, the existence and uniqueness of a solution of (90) in $\dot{H}^1(\mathbb{S}^{n-1})$ is proved.

Now, all solutions in $H^1(\mathbb{S}^{n-1})$ of (94) are of the form $\psi_{\eta\Sigma,V} + C_0$ where C_0 is any constant. Collecting all the solutions for all the possible $V \in \{\Omega_\Sigma\}^\perp$ leads to (93) and ends the proof. \square

Remark 8. We note that if Ω_Σ is changed into $-\Omega_\Sigma$, $\psi_{\eta\Sigma,V}$ must be changed into $\psi_{\eta\Sigma,-V}$. It follows that (93) remains unchanged.

We now define a vector-valued GCI $\vec{\psi}_{\eta\Sigma}$ in the following way

Definition 5.5. Given $(\eta, \Sigma) \in (0, \infty) \times \mathcal{U}_n^0$, we introduce the function $\vec{\psi}_{\eta\Sigma}: \mathbb{S}^{n-1} \rightarrow \mathbb{R}^n$, defined as the unique solution (in $\dot{H}^1(\mathbb{S}^{n-1})$) of the following vector-valued equation:

$$\nabla_\omega \cdot (G_{\eta\Sigma}(\omega) \nabla_\omega \vec{\psi}_{\eta\Sigma}) = (\omega \cdot \Omega_\Sigma) P_{\Omega_\Sigma^\perp} \omega G_{\eta\Sigma}(\omega), \quad \forall \omega \in \mathbb{S}^{n-1}.$$

We note that

$$\psi_{\eta\Sigma,V} = \vec{\psi}_{\eta\Sigma} \cdot V, \quad \forall V \in \{\Omega_\Sigma\}^\perp \quad \text{and} \quad \vec{\psi}_{\eta\Sigma} \cdot \Omega_\Sigma = 0,$$

and that $\vec{\psi}_{\eta\Sigma}$ is changed into $-\vec{\psi}_{\eta\Sigma}$ if Ω_Σ is changed into $-\Omega_\Sigma$.

We can provide an explicit expression of $\vec{\psi}_{\eta A_\Omega}$, for all $(\eta, \Omega) \in (0, \infty) \times \mathbb{S}^{n-1}$ as the next proposition shows. Let us first define the following space:

$$\mathcal{H} = \left\{ h : (-1, 1) \rightarrow \mathbb{R} \mid \int_{-1}^1 (1-r^2)^{\frac{n-1}{2}} |h(r)|^2 dr < \infty, \right. \\ \left. \int_{-1}^1 (1-r^2)^{\frac{n+1}{2}} |h'(r)|^2 dr < \infty \right\},$$

where h' denotes the derivative of h .

Proposition 9. Let $(\eta, \Omega) \in (0, \infty) \times \mathbb{S}^{n-1}$ be given. We have

$$\vec{\psi}_{\eta A_\Omega}(\omega) = h_\eta(\omega \cdot \Omega) \omega_\perp, \quad (96)$$

where $\omega_\perp = P_{\Omega^\perp} \omega$ and h_η is the unique solution in \mathcal{H} of the following equation:

$$-(1-r^2)^{\frac{n-1}{2}} e^{\eta r^2} (2\eta r^2 + n-1) h_\eta \\ + \frac{d}{dr} \left[(1-r^2)^{\frac{n+1}{2}} e^{\eta r^2} \frac{dh}{dr} \right] = r (1-r^2)^{\frac{n-1}{2}} e^{\eta r^2}. \quad (97)$$

Furthermore, h_η is odd and $h_\eta(r) \leq 0$ for $r \geq 0$.

Proof. We apply [21], Proposition 4.2 (ii) (with the following changes: $u \rightarrow \Omega$, $\frac{\kappa}{2} \rightarrow \eta$, $d \rightarrow n$, $\bar{\Gamma}^*(\psi, u) \rightarrow L_{\eta A_\Omega}^* \psi$). Note that these techniques were first developed in [19, 30]. \square

Remark 9. Formula (96) shows that the vector GCI $\vec{\psi}_{\eta A_\Omega}$ is invariant under rotations leaving Ω fixed. This is a consequence of the fact that A_Ω is uniaxial with axis Ω . No simple formula like (96) is available for more general vector GCI $\vec{\psi}_{\eta\Sigma}$, when $\Sigma \in \mathcal{U}_n^0$ is not uniaxial. However, while we will need vector GCI for general $\Sigma \in \mathcal{U}_n^0$, we will only need an explicit expression of them in the case of a uniaxial tensor $\Sigma = A_\Omega$. So, Prop. 9 is enough for our purpose.

The following proposition provides an alternate equation satisfied by h_η in terms of the function g defined in (53). Its proof is easy and is sketched in Appendix C.1 for the reader's convenience.

Proposition 10 (Alternate equation for h_η). *For $\theta \in [0, \pi]$, we define the function*

$$g(\theta) = -2\eta h_\eta(\cos \theta) \sin \theta. \tag{98}$$

Then g satisfies the equation (53).

Finally, the following proposition will have important consequences for the derivation of the macroscopic model:

Proposition 11. *Let $f: \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ be twice continuously differentiable such that $Q_f \neq 0$ and $\Sigma_f \in \mathcal{U}_n^0$. Then, the vector GCI $\vec{\psi}_{\eta_f \Sigma_f}$ is well-defined and we have*

$$\int_{\mathbb{S}^{n-1}} C(f) \vec{\psi}_{\eta_f \Sigma_f} d\omega = 0. \tag{99}$$

Remark 10. Proposition 11 expresses an important structural property of C . Let $(\eta, \Sigma) \in (0, \infty) \times \mathcal{U}_n^0$. The GCI $\vec{\psi}_{\eta \Sigma}$ cancels the collision operator acting on all functions f which satisfy $(\eta_f, \Sigma_f) = (\eta, \Sigma)$.

Proof. We show that $P_{\Omega_f^\perp}(Q_f \Omega_f) = 0$. Indeed, if this is the case, from (89), we get

$$\int_{\mathbb{S}^{n-1}} L_{\eta_f \Sigma_f} f \vec{\psi}_{\eta_f \Sigma_f} d\omega = 0,$$

and using (76), (87) and (88), this shows (99). But, by definition, Ω_f is the leading eigenvector of Q_f with eigenvalue λ_f . So, $Q_f \Omega_f = \lambda_f \Omega_f$ and thus $P_{\Omega_f^\perp}(Q_f \Omega_f) = 0$, which ends the proof. \square

Thanks to the GCI, we can now find how (83) translates into an equation for the Q-tensor principal direction Ω . This will be done below but first we provide some discussion of the GCI concept.

5.3. Discussion of the GCI concept.

5.3.1. *Rationale for Definition 89.* First, let us note that the condition $P_{\Omega_\Sigma^\perp}(Q_f \Omega_\Sigma) = 0$ involved in Definition 5.3 simply means that Ω_Σ is an eigenvector of Q_f . We now try to provide a geometric interpretation of Condition (89). First let us introduce a few additional notations. We endow \mathcal{S}_n^0 with the inner-product $S : P = \text{Tr}\{SP\}$ and for a subset B of \mathcal{S}_n^0 , its orthogonal with respect to this inner-product is denoted by B^\perp . We recall that B^\perp is a linear subspace of \mathcal{S}_n^0 and that $(B^\perp)^\perp = \text{Span}(B)$.

We now define the submanifold \mathcal{N} of \mathcal{U}_n^0 which consists of normalized prolate uniaxial Q-tensors i.e.

$$\mathcal{N} = \{A_\Omega \mid \Omega \in \mathbb{P}^{n-1}\} = \{\Omega \otimes \Omega - \frac{1}{n} \text{Id} \mid \Omega \in \mathbb{P}^{n-1}\}.$$

Note that \mathcal{N} is the manifold spanned by the NQT's of the equilibria (see Remark 7). The mapping $\mathbb{P}^{n-1} \ni \Omega \mapsto A_\Omega \in \mathcal{N}$ is a diffeomorphism. The tangent space of \mathcal{N} at A_Ω is given by:

$$T_{A_\Omega} \mathcal{N} = \{\Omega \otimes V + V \otimes \Omega \mid V \in \{\Omega\}^\perp\}. \tag{100}$$

Indeed, for $V \in T_\Omega \mathbb{P}^{n-1} = \{\Omega\}^\perp$, consider a curve $I \ni t \mapsto \xi(t) \in \mathbb{P}^{n-1}$ where I is an open interval of \mathbb{R} containing 0, such that $\xi(0) = \Omega$ and $\xi'(0) = V$. Then, $\frac{d}{dt}(A_{\xi(t)}) = \Omega \otimes V + V \otimes \Omega$, showing the claim. We denote by $P_{T_{A_\Omega} \mathcal{N}}$ the orthogonal projection of \mathcal{S}_n^0 on $T_{A_\Omega} \mathcal{N}$ for the inner product defined just above.

We have a mapping $p: \mathcal{U}_n^0 \rightarrow \mathcal{N}$, $\Sigma \mapsto A_{\Omega_\Sigma}$. For any $\Omega \in \mathbb{P}^{n-1}$, the pre-image $p^{-1}(\{A_\Omega\})$ is denoted by \mathcal{F}_Ω . All these pre-images are homeomorphic to

one-another. Let us choose one of them and denote it by \mathcal{F} . This endows \mathcal{U}_n^0 of a fiber bundle structure of base \mathcal{N} and fiber \mathcal{F} . Now, we have the following lemma:

Lemma 5.6. *Let $\Omega \in \mathbb{P}^{n-1}$ be given.*

- (i) *Let $Q \in \mathcal{S}_0^n$. Then, $P_{\Omega^\perp}(Q\Omega) = 0 \iff Q \in (T_{A_\Omega}\mathcal{N})^\perp$.*
- (ii) *\mathcal{F}_Ω is a subset of $(T_{A_\Omega}\mathcal{N})^\perp$.*

Proof. (i) Using the symmetry of Q and (100), we have:

$$\begin{aligned} P_{\Omega^\perp}(Q\Omega) = 0 &\iff (Q\Omega) \cdot V = 0, \quad \forall V \in \{\Omega\}^\perp \\ &\iff Q : (\Omega \otimes V) = 0, \quad \forall V \in \{\Omega\}^\perp \\ &\iff Q : (\Omega \otimes V + V \otimes \Omega) = 0, \quad \forall V \in \{\Omega\}^\perp \\ &\iff Q : B = 0, \quad \forall B \in T_{A_\Omega}\mathcal{N} \iff Q \in (T_{A_\Omega}\mathcal{N})^\perp, \end{aligned}$$

which shows (i).

(ii) Suppose $\Sigma \in \mathcal{F}_\Omega$. Then $A_{\Omega_\Sigma} = A_\Omega$ which implies $\Omega_\Sigma = \Omega$ (in \mathbb{P}^{n-1}). Thus, Ω is an eigenvector of Σ i.e. $P_{\Omega^\perp}(\Sigma\Omega) = 0$. Hence, by (i), $\Sigma \in (T_{A_\Omega}\mathcal{N})^\perp$. \square

So, Eq. (89) can be equivalently written:

$$\int_{\mathbb{S}^{n-1}} (L_{\eta\Sigma}f) \psi \, d\omega = 0 \quad \text{for all } f \text{ such that } Q_f \in (T_{A_{\Omega_\Sigma}}\mathcal{N})^\perp. \quad (101)$$

This can be geometrically interpreted as follows: to any $\Sigma \in \mathcal{U}_n^0$ we consider its projection (in the fiber bundle sense) $p(\Sigma) = A_{\Omega_\Sigma}$ onto \mathcal{N} . Then, (101) means that the GCI associated to (η, Σ) are all the functions ψ whose integrals against $L_{\eta\Sigma}f$ cancel when Q_f belongs to the orthogonal of the tangent space to \mathcal{N} at A_{Ω_Σ} . This is illustrated in Fig. 2. It is likely that this geometrical structure persists with other collision operators as it seems to express some intrinsic geometrical constraint. This point will be further developed in future work.

5.3.2. Relation between the GCI and the linearized collision operator. Let $D_f C$ the linearization of the collision operator C about the distribution function f and let $D_f C^*$ be its formal L^2 -adjoint. For a distribution function f , we call (η_f, Σ_f) the ‘moment’ of f . In this section, we show the following: suppose $(\eta, \Sigma) \in (0, \infty) \times \mathcal{U}_0^n$ is the moment of an equilibrium distribution function, i.e. $(\eta, \Sigma) = (\eta(\rho), A_\Omega)$ where $(\rho, \Omega) \in (\rho^*, \infty) \times \mathbb{S}^{n-1}/\{\pm 1\}$ and denote by $f^0 = \rho G_{\eta(\rho)A_\Omega}$ the corresponding equilibrium. Then, we have

$$\mathcal{C}_{\eta(\rho)A_\Omega} = \ker(D_{f^0} C^*). \quad (102)$$

On the other hand, if (η, Σ) is not the moment of an equilibrium, then, although there exist Gibbs distributions $f = \rho G_{\eta\Sigma}$ associated with (η, Σ) , in general, we have

$$\mathcal{C}_{\eta\Sigma} \neq \ker(D_f C^*). \quad (103)$$

Thus, a GCI associated to an arbitrary moment (η, Σ) is in general not in the kernel of the adjoint linearized collision operator about the corresponding Gibbs distribution. It is only so if (η, Σ) is the moment of an equilibrium in the above sense. Consequently, GCI are different and truly more general concepts than elements of such kernels. Likewise, Eq. (98) linking the GCI to the auxiliary function g given by (53) is only valid for moments $(\eta(\rho), A_\Omega)$ related to equilibria. Observe however that we will not need to explicit the form of the GCI for general moments, but only for those corresponding to an equilibrium (see Section 6 below).




FIGURE 2. Graphical representation of Condition (101). The ambient three-dimensional space in the figure represents the flat space \mathcal{S}_0^n in which \mathcal{U}_0^n is an imbedded manifold represented by a surface. \mathcal{N} is a submanifold of \mathcal{U}_0^n depicted as the curvy blue line. It endows \mathcal{U}_0^n of a fiber bundle structure of base \mathcal{N} . Let $\Sigma \in \mathcal{U}_0^n$. It projects (in the bundle sense) onto $A_\Omega \in \mathcal{N}$ and so, belongs to the fiber \mathcal{F}_Ω represented by the curvy red line. The tangent space to \mathcal{N} at A_Ω , $T_{A_\Omega}\mathcal{N}$ is represented by the magenta straight line. Its orthogonal $(T_{A_\Omega}\mathcal{N})^\perp$ is the gray-shaded plane on the figure. It contains \mathcal{F}_Ω by virtue of Lemma 5.6 (ii). Then, condition (101) means that the GCI associated with (η, Σ) are the functions ψ that cancel $L_{\eta\Sigma}f$ for all f whose Q-tensor Q_f (represented by the point Q on the figure) belongs to $(T_{A_\Omega}\mathcal{N})^\perp$.

Formula (102) is unsurprising. Indeed, Eq. (62) has been shown in [37, 59] using the Hilbert expansion method. This method corresponds to inserting the Hilbert expansion $f^\varepsilon = f^0 + \varepsilon f^1 + \mathcal{O}(\varepsilon^2)$ into the kinetic equation (40) and matching identical powers of ε . We get

$$C(f^0) = 0, \quad DC_{f^0}f^1 = T_{u^0}f^0,$$

for the terms of order ε^{-1} and ε^0 respectively (note that we also need to Hilbert-expand the velocity u^ε). Now, the first equation implies that f^0 is an equilibrium $f^0 = \rho G_{\eta(\rho)A_\Omega}$. Then, one looks for a necessary and sufficient condition for the existence of a solution f^1 to the second equation. Assuming that $\text{Im } DC_{f^0} = (\ker DC_{f^0}^*)^\perp$ (which can be proved via a careful study of the spectral properties of DC_{f^0} , see [59]), such a condition is

$$\int_{\mathbb{S}^{n-1}} T_{u^0}f^0 \psi \, d\omega = 0, \quad \forall \psi \in \ker DC_{f^0}^*.$$

Since this is also what we get when ψ ranges in $\mathcal{C}_{\eta(\rho)A_\Omega}$ (see Eq. (121) below), Eq. (102) must be true. However, it would be desirable to have a direct proof of (102). This is our goal here. As a by-product, we will also see why we have (103). We first compute the adjoint linearized collision operator.

Lemma 5.7 (Adjoint linearized collision operator). *Let $\rho \in (0, \infty)$, $S \in \mathcal{S}_0^n$. We have*

$$D_{\rho G_S} C^* g(\omega) = L_{\alpha \rho Q_{\rho G_S}}^* g(\omega) - \alpha \rho (\rho Q)_{G_S L_S^* g} : \omega \otimes \omega, \quad (104)$$

where G_S is defined by (43), the auxiliary operator L by (75) and L^* is its formal L^2 -adjoint. Here $(\rho Q)_{G_S L_S^* g}$ stands for the right-hand side of (10) with f replaced by $G_S L_S^* g$ (note that $\rho_{G_S L_S^* g} = 0$ so that $Q_{G_S L_S^* g}$ is not defined but $(\rho Q)_{G_S L_S^* g}$ itself is well-defined).

Proof. From (39) and the fact that U_f^0 depends linearly on f , we get

$$D_{\rho G_S} C f = \nabla_\omega \cdot (\nabla \omega f + f \nabla_\omega U_{\rho G_S}^0 + \rho G_S \nabla_\omega U_f^0). \quad (105)$$

We note that $\nabla_\omega U_{\rho G_S}^0 = -\nabla_\omega (\log G_{\alpha \rho Q_{\rho G_S}})$. Inserting this into (105), we get

$$D_{\rho G_S} C f = L_{\alpha \rho Q_{\rho G_S}} f + \rho G_S L_S^* U_f^0. \quad (106)$$

Thanks to (29), we also note that $L_S^* U_f^0 = L_S^* \check{U}_f^0$ with $\check{U}_f^0 = -\alpha(\omega \cdot \rho_f Q_f \omega)$. Thus, using (106), Stokes formula and that $L_S(G_S g) = G_S L_S^* g$, we get

$$\int_{\mathbb{S}^{n-1}} D_{\rho G_S} C f g d\omega = \int_{\mathbb{S}^{n-1}} f L_{\alpha \rho Q_{\rho G_S}}^* g d\omega + \rho \int_{\mathbb{S}^{n-1}} \check{U}_f^0 G_S L_S^* g d\omega.$$

Inserting the expression of \check{U}_f^0 into this formula, using the expression (10) of $\rho_f Q_f$ and exchanging ω and ω' in the resulting integral, we are led to (104). \square

Now, in the case of an equilibrium, we compute the kernel of the adjoint linearized collision operator:

Lemma 5.8 (kernel of $D_{f^0} C^*$ when f^0 is an equilibrium). *Let $\rho \in (\rho^*, \infty)$ and $\Omega \in \mathbb{S}^{n-1}/\{\pm 1\}$. Let $f^0 = \rho G_{\eta(\rho)A_\Omega}$ be an equilibrium of C , where the function $\rho \mapsto \eta(\rho)$ is defined in Prop. 3. Define $\tilde{\mathcal{X}}_{\rho, \Omega}$ to be the space of functions $\varphi : \omega \mapsto \varphi(\omega)$ which satisfy*

$$\varphi(\omega) = \alpha \rho (\rho Q)_{G_{\eta(\rho)A_\Omega} \varphi} : \omega \otimes \omega, \quad \forall \omega \in \mathbb{S}^{n-1}. \quad (107)$$

Then we have

$$g \in \ker(D_{\rho G_{\eta(\rho)A_\Omega}} C^*) \iff \int_{\mathbb{S}^{n-1}} L_{\eta(\rho)A_\Omega} f g d\omega = 0, \quad \forall f \in \tilde{\mathcal{X}}_{\rho, \Omega}^\perp, \quad (108)$$

where the orthogonality is with respect to the standard $L^2(\mathbb{S}^{n-1})$ -inner product.

Proof. Defining $S = \eta(\rho)A_\Omega$, we have

$$\alpha \rho Q_{\rho G_S} = \alpha \rho Q_{\rho G_{\eta(\rho)A_\Omega}} = \eta A_\Omega = S, \quad (109)$$

thanks to (48) and (49). Thus, thanks to (104), we are led to

$$D_{\rho G} C^* g(\omega) = L^* g(\omega) - \alpha \rho (\rho Q)_{G L^* g} : \omega \otimes \omega, \quad (110)$$

where here and in the remainder of the proof, we omit the dependence of η on ρ , as well as the index ηA_Ω on L^* and G and the indices ρ, Ω on $\tilde{\mathcal{X}}$ for clarity.

For any smooth enough function f , we have by the Stokes formula:

$$\int_{\mathbb{S}^{n-1}} L f g d\omega = \int_{\mathbb{S}^{n-1}} f L^* g d\omega = \int_{\mathbb{S}^{n-1}} f \varphi d\omega,$$

with $\varphi = L^* g$. Thanks to (110) and the fact that $g \in \ker D_{\rho G} C^*$, φ satisfies (107), so $\varphi \in \tilde{\mathcal{X}}$. If $f \in \tilde{\mathcal{X}}^\perp$, we deduce that $\int L f g d\omega = 0$, which shows the left-to-right implication of (108).

Conversely suppose that g is such that $\int Lf g d\omega = 0, \forall f \in \tilde{\mathcal{X}}^\perp$, i.e.

$$f \in \tilde{\mathcal{X}}^\perp \implies f \in \{L^*g\}^\perp.$$

Taking the orthogonals, we get

$$\text{Span}\{L^*g\} \subset \tilde{\mathcal{X}}.$$

Indeed, both $\text{Span}\{L^*g\}$ and $\tilde{\mathcal{X}}$ are finite-dimensional, hence closed. This is obvious for the former which is one-dimensional. For the latter, by (107), $\tilde{\mathcal{X}}$ is included in the space of quadratic polynomials in ω , which is a finite-dimensional space. So, defining $\varphi = L^*g$, we have $\varphi \in \tilde{\mathcal{X}}$. Replacing φ by its expression in terms of g in (107), we get $D_{\rho G}C^*(g) = 0$, which shows the right-to-left implication of (108) and ends the proof. \square

Next, we prove an alternate characterization of the space $\tilde{\mathcal{X}}_{\rho, \Omega}$.

Lemma 5.9. *Let ρ, Ω, f^0 and η as in Lemma 5.8. Then,*

$$\tilde{\mathcal{X}}_{\rho, \Omega} = \mathcal{X}_\Omega, \quad (111)$$

where \mathcal{X}_Ω is defined by (91).

Proof. Let $\varphi \in \tilde{\mathcal{X}}$ (using the simplified notations of the previous proof). From (107), we have $\varphi(\omega) = K : \omega \otimes \omega$ where $K = \alpha\rho(\rho Q)_{G\varphi}$. Hence, K satisfies the fixed point equation

$$K = \alpha\rho(\rho Q)_{GK:\omega \otimes \omega}, \quad (112)$$

which implies that

$$\text{Tr}K = 0. \quad (113)$$

Using (10), (20) and (113), we can develop (112) into:

$$K = \alpha\rho \mathbb{T}_{G\eta A_\Omega} : K. \quad (114)$$

According to (156), there are three real numbers $a_k, k = 1, \dots, 3$, such that

$$\mathbb{T}_{G\eta A_\Omega} = a_1 \Omega^{\otimes 4} + 6a_2 (\Omega \otimes \Omega \otimes \text{Id})_s + 3a_3 (\text{Id} \otimes \text{Id})_s. \quad (115)$$

We uniquely define $V \in \{\Omega\}^\perp$ and $r \in \mathbb{R}$ by $K\Omega = r\Omega + V$. inserting (115) into (114) and using (113), we get

$$\left(\frac{1}{2\alpha\rho} - a_3\right)K = a_2 (\Omega \otimes V + V \otimes \Omega) + \frac{1}{2}((a_1 + 4a_2)r \Omega \otimes \Omega + a_2 r \text{Id}) \quad (116)$$

We now state the following lemma, whose proof can be found in Appendix C.2

Lemma 5.10. *We have*

$$\frac{1}{2\alpha\rho} - a_3 = a_2 \neq 0, \quad (117)$$

$$a_1 + (n+4)a_2 = S_2(\eta). \quad (118)$$

Using (117), Eq. (116) leads to

$$K = \Omega \otimes V + V \otimes \Omega + \frac{1}{2a_2} [(a_1 + 4a_2)r \Omega \otimes \Omega + a_2 r \text{Id}].$$

With (113), we get

$$0 = \text{Tr}K = \frac{1}{2a_2} [(a_1 + (n+4)a_2)] r,$$

which, with (118) and the fact that $S_2(\eta) \neq 0$ (see Prop. 2 (iii)), leads to $r = 0$ and

$$K = \Omega \otimes V + V \otimes \Omega.$$

Thus,

$$\varphi = 2(\Omega \cdot \omega)(V \cdot \omega). \quad (119)$$

Reciprocally, by similar but simpler computations, we easily get that φ given by (119) with arbitrary $V \in \{\Omega\}^\perp$ satisfies (112). In the end, we find

$$\tilde{\mathcal{X}} = \{(\Omega \cdot \omega)(V \cdot \omega) \mid V \in \{\Omega\}^\perp\} = \mathcal{X}_\Omega,$$

which ends the proof. \square

We can now state the following

Theorem 5.11. *Let $f^0 = \rho G_{\eta(\rho)A_\Omega}$ be an equilibrium of C . Then, we have*

$$\mathcal{C}_{\eta(\rho)A_\Omega} = \ker(D_{\rho G_{\eta(\rho)A_\Omega}} C^*),$$

where $\mathcal{C}_{\eta(\rho)A_\Omega}$ is the space of GCI associated with the equilibrium moments $(\eta(\rho), A_\Omega)$ (see Definition 5.3).

Proof. Indeed, we have the sequence of equivalences:

$$\psi \text{ is a GCI associated with } (\eta(\rho), A_\Omega) \iff$$

$$\iff \left(f \in \mathcal{X}_\Omega^\perp \implies \int_{\mathbb{S}^{n-1}} L_{\eta(\rho)A_\Omega} f \psi \, d\omega = 0 \right)$$

$$\iff \left(f \in \tilde{\mathcal{X}}_{\rho, \Omega}^\perp \implies \int_{\mathbb{S}^{n-1}} L_{\eta(\rho)A_\Omega} f \psi \, d\omega = 0 \right)$$

$$\iff \psi \in \ker(D_{\rho G_{\eta(\rho)A_\Omega}} C^*),$$

where the first equivalence comes from (89) and (92), the second one from (111) and the third one, from (108). This ends the proof. \square

The key property which led to Theorem 5.11 in the case where f^0 is an equilibrium is (109). It gave rise to the structure

$$D_{\rho G} C^* g = \varphi(\omega) - \alpha \rho (\rho Q)_{G_\varphi} : \omega \otimes \omega, \quad (120)$$

with $\varphi = L^* g$ which led to the definition of the space $\tilde{\mathcal{X}}_{\rho, \Omega}$. Now, if (η, Σ) is not a moment of an equilibrium, we have $\alpha \rho Q_{\rho G_S} \neq S$ as the equality is a characterization of the moments of equilibria. Then, by inspection of (104), we see that the structure (120) is lost and the proof cannot be continued. These considerations strongly support (103). Indeed, we have the following counter-example in dimension $n = 3$ whose proof can be found in Appendix C.3.

Proposition 12. *Let $n = 3$. Let $f = \rho G_{\eta A_\Omega}$ where $\eta \neq \eta(\rho)$ (in other words, in spite of being a Gibbs distribution, f is not an equilibrium). Then we have (103) (with $\Sigma = A_\Omega$).*

So, the space of GCI $\mathcal{C}_{\eta\Sigma}$ is related to important structural properties of C such as Prop. 99. By contrast, the space $\ker(D_f C^*)$ does not play any particular role. The exception is when the Gibbs distribution $\rho G_{\eta\Sigma}$ is an equilibrium, in which case the two spaces are equal. This shows that GCI are a more relevant and general concept than the space $\ker(D_f C^*)$ which appears in the Hilbert method.

6. Equation for the Q-tensor axis direction Ω .

6.1. Abstract derivation. In this section, we provide an abstract set of equations allowing us to determine the evolution equation for the Q-tensor axis direction Ω . We recall the expression (37) of $T_u(f)$. We have the:

Proposition 13. *Let $f = \lim_{\varepsilon \rightarrow 0} f^\varepsilon$ with $f(x, \omega, t) = \rho(x, t) G_{\eta(\rho(x, t))A_{\Omega(x, t)}}(\omega)$ for all $(x, t) \in \mathcal{B}$ where \mathcal{B} is given by (59) and the function $\rho \mapsto \eta(\rho)$ is defined in Prop. 3. Then, we have*

$$\int_{\mathbb{S}^{n-1}} T_u(\rho(x, t) G_{\eta(\rho(x, t))A_{\Omega(x, t)}}) \vec{\psi}_{\eta(\rho(x, t))A_{\Omega(x, t)}}(\omega) d\omega = 0, \quad (121)$$

where $\vec{\psi}_{\eta(\rho(x, t))A_{\Omega(x, t)}}$ is the vector GCI associated with $(\eta(\rho(x, t)), A_{\Omega(x, t)})$ (see Section 5.2).

Remark 11. We note that (121) is unchanged if $\Omega(x, t)$, and consequently $\vec{\psi}_{A_{\Omega(x, t)}}$, are changed in their opposites.

Proof. Let $(x, t) \in \mathcal{B}$ be given. For simplicity, in the proof, we omit the variables (x, t) . We also denote $\rho^\varepsilon := \rho_{f^\varepsilon}$, $Q^\varepsilon := Q_{f^\varepsilon}$, $\lambda^\varepsilon := \lambda_{f^\varepsilon}$, etc. and $\rho := \rho_f$, $Q := Q_f$, $\lambda := \lambda_f$, etc. By the fact that $f^\varepsilon \rightarrow f$, we get $\rho^\varepsilon Q^\varepsilon \rightarrow \rho Q = \rho \frac{n}{n-1} \lambda A_\Omega$, with $\frac{n}{n-1} \lambda = \frac{\eta(\rho)}{\alpha \rho}$. Since $\rho \neq 0$ (because $(x, t) \in \mathcal{B}$) and λ is a simple eigenvalue of Q , then, for ε small enough, $\rho^\varepsilon \neq 0$, $Q^\varepsilon \rightarrow Q$ and λ^ε is a simple eigenvalue of Q^ε such that $\lambda^\varepsilon \rightarrow \lambda$ (because the subset of \mathcal{S}_n^0 of matrices which have simple leading eigenvalue is an open set). Thus, $\Sigma^\varepsilon = \frac{n-1}{n \lambda^\varepsilon} Q^\varepsilon$ is defined, belongs to \mathcal{U}_n^0 and is such that $\Sigma^\varepsilon \rightarrow \Sigma = A_\Omega$ as $\varepsilon \rightarrow 0$.

By the smoothness of Σ^ε with respect to ε , we can find a smooth lifting of $\Omega_{\Sigma^\varepsilon} \in \mathbb{P}^{n-1}$ into $\Omega^\varepsilon \in \mathbb{S}^{n-1}$. Thus, we can form the GCI $\vec{\psi}_{\eta^\varepsilon \Sigma^\varepsilon}$ using this smooth determination of $\Omega_{\Sigma^\varepsilon}$ (remember that we need to fix the sign of $\Omega_{\Sigma^\varepsilon}$ because the sign of $\vec{\psi}_{\eta^\varepsilon \Sigma^\varepsilon}$ depends on it). This makes $\vec{\psi}_{\eta^\varepsilon \Sigma^\varepsilon}$ a smooth function of ε (because $\vec{\psi}_{\eta S}$ is a smooth function of $(\eta, S) \in [0, \infty) \times \mathcal{U}_0^n$) such that $\vec{\psi}_{\eta^\varepsilon \Sigma^\varepsilon} \rightarrow \vec{\psi}_{\eta A_\Omega}$ when $\varepsilon \rightarrow 0$.

Thanks to (99), we have

$$\int_{\mathbb{S}^{n-1}} C(f^\varepsilon) \vec{\psi}_{\eta^\varepsilon \Sigma^\varepsilon} d\omega = 0.$$

So, multiplying (40) by $\vec{\psi}_{\eta^\varepsilon \Sigma^\varepsilon}$, integrating the resulting expression with respect to ω leads to

$$\int_{\mathbb{S}^{n-1}} T_{u^\varepsilon}(f^\varepsilon) \vec{\psi}_{\eta^\varepsilon \Sigma^\varepsilon} d\omega = 0.$$

Now letting $\varepsilon \rightarrow 0$, with $u^\varepsilon \rightarrow u$, $f^\varepsilon \rightarrow \rho G_{\eta(\rho)A_\Omega}$, $\eta^\varepsilon \rightarrow \eta(\rho)$, $\Sigma^\varepsilon \rightarrow A_\Omega$, $\vec{\psi}_{\eta^\varepsilon \Sigma^\varepsilon} \rightarrow \vec{\psi}_{\eta(\rho)A_\Omega}$, we get (121). This ends the proof. \square

6.2. Derivation of the equation for Ω . In this section, we derive the explicit equation for Ω by inserting expression (96) into the abstract formulation (121) and compute the integral explicitly. This is summarized in the following

Proposition 14. *Let $f = \lim_{\varepsilon \rightarrow 0} f^\varepsilon = \rho(x, t) G_{\eta(\rho(x, t))A_{\Omega(x, t)}}$ as given in Corollary 1. Then, Ω satisfies (62)*

Proof of Proposition 14. For simplicity, we omit the dependencies of η and λ on ρ , of h_η on $(\omega \cdot \Omega)$, of $G_{\eta A_\Omega}$ on ω and of ρ and Ω on (x, t) . Inserting (96) into (121), we get:

$$V_\Omega := \int_{\mathbb{S}^{n-1}} T_u(\rho G_{\eta A_\Omega}) h_\eta \omega_\perp d\omega = 0. \quad (122)$$

We define

$$\begin{aligned} D_t &= \partial_t + u \cdot \nabla_x, \\ Af &= \nabla_\omega \cdot (f(\Lambda P_{\omega^\perp} E - W)\omega), \\ Bf &= 2\alpha\beta \nabla_\omega \cdot (f P_{\omega^\perp} \Delta_x(\rho_f Q_f)\omega), \end{aligned}$$

so that $T_u(f) = D_t f + Af + Bf$ and

$$V_\Omega = \int_{\mathbb{S}^{n-1}} (D_t + A + B)(\rho G_{\eta A_\Omega}) h_\eta \omega_\perp d\omega = V_\Omega^{(1)} + V_\Omega^{(2)} + V_\Omega^{(3)}. \quad (123)$$

Using (86) which gives $D_t \rho = 0$ and $D_t \eta = \eta' D_t \rho = 0$, where η' is the derivative of η with respect to ρ , we get

$$D_t(\rho G_{\eta A_\Omega}) = \rho G_{\eta A_\Omega} 2\eta(\omega \cdot \Omega)(P_{\Omega^\perp} \omega) \cdot D_t \Omega,$$

where we have used that the denominator of (45) does not depend on Ω . Then, we apply (141) and the fact that $D_t \Omega$ is orthogonal to Ω and get

$$V_\Omega^{(1)} = \tilde{\gamma}_1 D_t \Omega, \quad (124)$$

with

$$\tilde{\gamma}_1 = \frac{2\eta\rho}{n-1} \int_{\mathbb{S}^{n-1}} G_{\eta A_\Omega} h_\eta(\omega \cdot \Omega)(1 - (\omega \cdot \Omega)^2) d\omega. \quad (125)$$

Next, we have

$$\begin{aligned} A(\rho G_{\eta A_\Omega}) &= \nabla_\omega \cdot (\rho G_{\eta A_\Omega} (\Lambda P_{\omega^\perp} E - W)\omega) \\ &= \rho G_{\eta A_\Omega} [\nabla_\omega(\log G_{\eta A_\Omega}) \cdot (\Lambda P_{\omega^\perp} E - W)\omega + \nabla_\omega \cdot ((\Lambda P_{\omega^\perp} E - W)\omega)]. \end{aligned}$$

First, we compute $\nabla_\omega \cdot ((\Lambda P_{\omega^\perp} E - W)\omega)$. Let $X = \Lambda E - W$ for simplicity and let $(e_i)_{i=1, \dots, n}$ be the canonical basis of \mathbb{R}^n . Define $\mathcal{X}_i = \sum_{j=1}^n X_{ij} e_j$. Then, we can write $X = \sum_{i=1}^n e_i \otimes \mathcal{X}_i$. Then, $P_{\omega^\perp} X \omega = \sum_{i=1}^n (\mathcal{X}_i \cdot \omega) P_{\omega^\perp} e_i$. We note that $\nabla_\omega \cdot P_{\omega^\perp} e_i = \Delta_\omega(\omega \cdot e_i) = -(n-1)(\omega \cdot e_i)$ because $(\omega \cdot e_i)$ is a spherical harmonic of degree 1 hence an eigenfunction of the spherical laplacian associated to the eigenvalue $-(n-1)$. Thus,

$$\begin{aligned} \nabla_\omega \cdot (P_{\omega^\perp} X \omega) &= \sum_{i=1}^n \left[P_{\omega^\perp} \mathcal{X}_i \cdot P_{\omega^\perp} e_i - (n-1)(\mathcal{X}_i \cdot \omega)(\omega \cdot e_i) \right] \\ &= \sum_{i,j=1}^n X_{ij} (P_{\omega^\perp} e_i \cdot P_{\omega^\perp} e_j - (n-1)\omega_i \omega_j) = \sum_{i,j=1}^n X_{ij} (\delta_{ij} - n\omega_i \omega_j) \\ &= \text{Tr} X - nX : (\omega \otimes \omega), \end{aligned}$$

where δ_{ij} is the Kronecker symbol and $\text{Tr} X$ is the trace of X . Now, with $X = \Lambda E - W$, owing to the facts that $\text{Tr} X = \Lambda \nabla_x \cdot u = 0$ and remembering that E is symmetric and W , antisymmetric, we get

$$\begin{aligned} A(\rho G_{\eta A_\Omega}) &= \rho G_{\eta A_\Omega} [2\eta(\omega \cdot \Omega) P_{\omega^\perp} \Omega \cdot (\Lambda P_{\omega^\perp} E - W)\omega - n\Lambda E : (\omega \otimes \omega)] \\ &= \rho G_{\eta A_\Omega} \left\{ \Lambda [2\eta(\omega \cdot \Omega) P_{\omega^\perp} \Omega \otimes \omega - n\omega \otimes \omega] : E - 2\eta(\omega \cdot \Omega) (P_{\omega^\perp} \Omega \otimes \omega) : W \right\}. \end{aligned}$$

Using the decomposition (142), we get $P_{\omega^\perp}\Omega = (1 - (\omega \cdot \Omega)^2)\Omega - (\omega \cdot \Omega)\omega_\perp$, and so,

$$\begin{aligned} & A(\rho G_{\eta A_\Omega}) \\ &= \rho G_{\eta A_\Omega} 2\eta(\omega \cdot \Omega) \left[\Lambda \left(1 - \frac{n}{\eta} - 2(\omega \cdot \Omega)^2 \right) (\omega_\perp \otimes \Omega) : E + (\omega_\perp \otimes \Omega) : W \right] \\ & \qquad \qquad \qquad + \text{even tensor powers of } \omega_\perp. \end{aligned}$$

Now, multiplying by $h\omega_\perp$ and integrating over ω , the resulting odd tensor powers of ω_\perp vanish in the integration thanks to (140). Thanks to (141), we find that

$$V_\Omega^{(2)} = \tilde{\gamma}_1 W\Omega + \tilde{\gamma}_3 \Lambda P_{\Omega^\perp} E\Omega, \tag{126}$$

with

$$\tilde{\gamma}_3 = \left(1 - \frac{n}{\eta} \right) \tilde{\gamma}_1 - 2\tilde{\gamma}_2, \tag{127}$$

$$\tilde{\gamma}_2 = \frac{2\eta\rho}{n-1} \int_{\mathbb{S}^{n-1}} G_{\eta A_\Omega} h_\eta (\omega \cdot \Omega)^3 (1 - (\omega \cdot \Omega)^2) d\omega. \tag{128}$$

The computation of $V_\Omega^{(3)}$ is the same as that of $V_\Omega^{(2)}$ with $\Lambda E - W$ replaced by $2\alpha\beta \Delta_x(\rho Q_{G_{\eta A_\Omega}})$. Since $\Delta_x(\rho Q_{G_{\eta A_\Omega}})$ is a symmetric trace-free tensor, we get from (126):

$$V_\Omega^{(3)} = 2\alpha\beta\tilde{\gamma}_3 P_{\Omega^\perp} (\Delta_x(\rho Q_{G_{\eta A_\Omega}})\Omega).$$

With (50), we get

$$\begin{aligned} \alpha\Delta_x(\rho Q_{G_{\eta A_\Omega}}) &= \Delta_x(\eta A_\Omega) = \Delta_x\eta A_\Omega + 4 \left[\left((\nabla_x\eta \cdot \nabla_x)\Omega \right) \otimes \Omega \right]_s \\ & \qquad \qquad \qquad + 2\eta \left((\nabla_x\Omega)^T (\nabla_x\Omega) + [(\Delta_x\Omega) \otimes \Omega]_s \right), \end{aligned} \tag{129}$$

where the index s means the symmetric part of a tensor (i.e. $S_s = \frac{1}{2}(S + S^T)$ for an $n \times n$ matrix S). Then, owing to the fact that any derivative of Ω is orthogonal to Ω , we have

$$\alpha\Delta_x(\rho Q_{G_{\eta A_\Omega}})\Omega = \Delta_x\eta\Omega + 2(\nabla_x\eta \cdot \nabla_x)\Omega + \eta(\Delta_x\Omega + (\Omega \cdot \Delta_x\Omega)\Omega),$$

and with (82),

$$\alpha P_{\Omega^\perp} (\Delta_x(\rho Q_{G_{\eta A_\Omega}})\Omega) = 2(\nabla_x\eta \cdot \nabla_x)\Omega + \eta P_{\Omega^\perp} \Delta_x\Omega = P_{\Omega^\perp} \Delta_x(\eta\Omega).$$

It follows that

$$V_\Omega^{(3)} = 2\beta\tilde{\gamma}_3 P_{\Omega^\perp} \Delta_x(\eta\Omega). \tag{130}$$

Inserting (124), (126), (130), into (123), we get

$$V_\Omega = \tilde{\gamma}_1 (D_t\Omega + W\Omega) + \tilde{\gamma}_3 P_{\Omega^\perp} (\Lambda E\Omega + 2\beta \Delta_x(\eta\Omega)).$$

So, with (122) and (127), we get (62) with

$$c = -\Lambda \frac{\tilde{\gamma}_3}{\tilde{\gamma}_1} = \Lambda \left(\frac{n}{\eta} - 1 + 2\frac{\tilde{\gamma}_2}{\tilde{\gamma}_1} \right). \tag{131}$$

Now, the following formulas are shown in the Appendix D.1:

$$\tilde{\gamma}_3 = \frac{\rho S_2(\eta)}{2\eta}, \quad \tilde{\gamma}_1 = -\frac{\rho}{2\eta(n-1)} \langle\langle g \frac{d\tilde{U}_0}{d\theta} \rangle\rangle_{e^{\eta \cos^2 \theta}}. \tag{132}$$

Thus, (131) leads to (54) and ends the proof. \square

We now investigate under which conditions $\frac{c}{\Lambda}$ is non-negative:

Proof of Proposition 4. From Prop. 2 (iii), we know that the $(n-1)S_2(\eta) > 0$. Now, Prop. 9 and Eqs. (98) and (52) show that both $g(\theta)$ and $\frac{d\tilde{U}_0}{d\theta}(\theta)$ have the sign

of $\cos \theta$. This implies that $g(\theta) \frac{d\tilde{U}_\omega}{d\theta}(\theta)$ is positive on $[0, \pi]$ and consequently, that the denominator of (54) is positive. Altogether, this shows that $\frac{c}{\Lambda} > 0$ and ends the proof. \square

7. Conclusion. We have investigated the passage from the Doi-Navier-Stokes model of liquid crystals to the Ericksen-Leslie system when the Deborah number goes to zero. By contrast to previous literature, we have developed a moment method, exploiting the conservations satisfied by the collision operator. These conservations are of a non-classical type and have required the development of a new concept, the generalized collision invariants. Their link to geometrical and analytical structures of the collision operator has been discussed and their use for the derivation of the limit model has been detailed. This derivation has been achieved in arbitrary dimensions and assuming a full spatio-temporal dependence of the polymer molecule density. The latter generates additional terms in the Ericksen stresses that have not been previously described in the literature.

This work opens many research directions. The first one is the development of a rigorous convergence result using this moment method. This is a quite challenging task but one may hope that, if successful, it would lead to a result in a weaker setting than the currently available results. The energetic properties of the limit model must be investigated. A proof that the extra terms appearing in the Oseen-Franck energy due to the spatio-temporal dependence of the polymer molecule density lead to a positive energy is missing at the present time. This would be a necessary step for a well-posedness theory of the resulting Ericksen-Leslie system. In spite of using Q-tensors as auxiliary quantities, the Doi model and its limit, the Ericksen-Leslie system are, in essence, vector models, i.e. models for polymer orientations only. Currently, attempts are being made to build truly tensorial models in association with Landau-de Gennes energies i.e. energies depending on the local average Q-tensor and its gradients. This is clearly an interesting playground to test the applicability of the GCI concept to more general situations.

Appendix A. Appendix to Section 2 on Doi's model.

A.1. **Proof of the virtual work principle (16).** We have, with (15):

$$\begin{aligned} \frac{d\mathcal{A}^R}{dt} &= \left\langle \frac{\delta \mathcal{A}^R}{\delta f}(f), \frac{\partial f}{\partial t} \right\rangle = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} \mu_f^R \frac{\partial f}{\partial t} dx d\omega \\ &= \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} \mu_f^R \left\{ -\nabla_x \cdot (u f) - \nabla_\omega \cdot (f (\Lambda P_{\omega^\perp} E - W)\omega) \right. \\ &\quad \left. + \frac{D}{k_B T} \nabla_\omega \cdot (f \nabla_\omega \mu_f^R) \right\} dx d\omega \\ &=: \text{I} + \text{II} + \text{III} \end{aligned}$$

Using Stokes's formula, assuming that all terms vanish at infinity and with (17), we find

$$\text{I} = - \int_{\mathbb{R}^n} F_f^R \cdot u dx, \quad \text{III} = - \frac{D}{k_B T} \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} f |\nabla_\omega \mu_f^R|^2 dx d\omega.$$

Then, using Stokes's formula, the fact that $\nabla_\omega \mu_f^R \cdot \omega = 0$ and straightforward tensor algebra, we have

$$\text{II} = \int_{\mathbb{R}^n \times \mathbb{S}^{n-1}} f (\Lambda E - W)\omega \cdot \nabla_\omega \mu_f^R dx d\omega$$

$$\begin{aligned}
 &= \int_{\mathbb{R}^n} \left(\int_{\mathbb{S}^{n-1}} f(\omega \otimes \nabla_\omega \mu_f^R) d\omega \right) : (\Lambda E + W) dx \\
 &= \int_{\mathbb{R}^n} \left(\int_{\mathbb{S}^{n-1}} f \left[\frac{\Lambda + 1}{2} (\omega \otimes \nabla_\omega \mu_f^R) + \frac{\Lambda - 1}{2} (\nabla_\omega \mu_f^R \otimes \omega) \right] d\omega \right) : \nabla_x u dx \\
 &= \int_{\mathbb{R}^n} \left(\int_{\mathbb{S}^{n-1}} f \left[\Lambda (\omega \otimes \nabla_\omega \mu_f^R)_s - (\omega \otimes \nabla_\omega \mu_f^R)_a \right] d\omega \right) : \nabla_x u dx \\
 &= \int_{\mathbb{R}^n} \sigma_f^R : \nabla_x u dx.
 \end{aligned}$$

This leads to (16). \square

A.2. Proofs of Formulas (21) and (22) for the extra-stresses. We begin with a Lemma:

Lemma A.1. *Let f and $\varphi: \mathbb{S}^{n-1} \rightarrow \mathbb{R}$ be two smooth functions. Then, we have*

$$\int_{\mathbb{S}^{n-1}} \nabla_\omega f \varphi d\omega = - \int_{\mathbb{S}^{n-1}} f \nabla_\omega \varphi d\omega + (n-1) \int_{\mathbb{S}^{n-1}} f \varphi \omega d\omega. \quad (133)$$

Proof. Let $B \in \mathbb{R}^n$ be a fixed vector and denote by X the left-hand side of (133). Then, using Stokes formula, we have

$$\begin{aligned}
 X \cdot B &= \int_{\mathbb{S}^{n-1}} \nabla_\omega f \cdot B \varphi d\omega = \int_{\mathbb{S}^{n-1}} \nabla_\omega f \cdot P_{\omega^\perp} B \varphi d\omega \\
 &= - \int_{\mathbb{S}^{n-1}} f \nabla_\omega \cdot (P_{\omega^\perp} B \varphi) d\omega \\
 &= - \int_{\mathbb{S}^{n-1}} f \nabla_\omega \cdot (P_{\omega^\perp} B) \varphi d\omega - \int_{\mathbb{S}^{n-1}} f P_{\omega^\perp} B \cdot \nabla_\omega \varphi d\omega.
 \end{aligned}$$

We have

$$\nabla_\omega \cdot (P_{\omega^\perp} B) = \nabla_\omega \cdot \nabla_\omega (\omega \cdot B) = \Delta_\omega (\omega \cdot B) = -(n-1) \omega \cdot B,$$

where the last identity follows from the fact that the function $\omega \mapsto \omega \cdot B$ is a spherical harmonic of degree 1. Thus,

$$X \cdot B = (n-1) \int_{\mathbb{S}^{n-1}} f \varphi \omega \cdot B d\omega - \int_{\mathbb{S}^{n-1}} f \nabla_\omega \varphi \cdot B d\omega,$$

which leads to (133). \square

Proof of (21). Inserting (13) into the first equation of (17), we have $\sigma_f^R = \Lambda \bar{\sigma}_s + \bar{\sigma}_a$ with

$$\begin{aligned}
 \bar{\sigma} &= \int_{\mathbb{S}^{n-1}} f(\omega \otimes \nabla_\omega \mu_f^R) d\omega \\
 &= k_B T \int_{\mathbb{S}^{n-1}} \omega \otimes \nabla_\omega f d\omega + \int_{\mathbb{S}^{n-1}} f(\omega \otimes \nabla_\omega U_f^R) d\omega. \quad (134)
 \end{aligned}$$

Using (133) with $\varphi = \omega_i$, we get

$$\begin{aligned}
 \int_{\mathbb{S}^{n-1}} \omega_i \nabla_\omega f d\omega &= - \int_{\mathbb{S}^{n-1}} \nabla_\omega \omega_i f d\omega + (n-1) \int_{\mathbb{S}^{n-1}} f \omega \omega_i d\omega \\
 &= - \int_{\mathbb{S}^{n-1}} P_{\omega^\perp} e_i f d\omega + (n-1) \int_{\mathbb{S}^{n-1}} f \omega (\omega \cdot e_i) d\omega
 \end{aligned}$$

$$\begin{aligned}
&= - \int_{\mathbb{S}^{n-1}} (e_i - (e_i \cdot \omega)\omega) f \, d\omega + (n-1) \int_{\mathbb{S}^{n-1}} f (e_i \cdot \omega)\omega \, d\omega \\
&= n \int_{\mathbb{S}^{n-1}} f (\omega(\omega \cdot e_i) - \frac{1}{n}e_i) \, d\omega,
\end{aligned}$$

where e_i denotes the i -th vector of the canonical basis of \mathbb{R}^n . In view of (10), it follows that $\int_{\mathbb{S}^{n-1}} \omega \otimes \nabla_\omega f \, d\omega = n\rho_f Q_f$. Inserting this in (134) leads to

$$\bar{\sigma} = nk_B T \rho_f Q_f + \int_{\mathbb{S}^{n-1}} f (\omega \otimes \nabla_\omega U_f^R) \, d\omega,$$

which, in turn, leads to (21). \square

Proof of (22). We multiply Doi's equation (15) by $\omega \otimes \omega - \frac{1}{n}\text{Id}$ and integrate it with respect to ω . This leads to

$$\begin{aligned}
0 &= \int_{\mathbb{S}^{n-1}} (\partial_t f + \nabla_x \cdot (uf)) (\omega \otimes \omega - \frac{1}{n}\text{Id}) \, d\omega \\
&+ \int_{\mathbb{S}^{n-1}} \nabla_\omega \cdot (f(\Lambda P_{\omega^\perp} E - W)\omega) (\omega \otimes \omega - \frac{1}{n}\text{Id}) \, d\omega \\
&- \frac{D}{k_B T} \int_{\mathbb{S}^{n-1}} \nabla_\omega \cdot (f \nabla_\omega \mu_f^R) (\omega \otimes \omega - \frac{1}{n}\text{Id}) \, d\omega \\
&=: \text{I} + \text{II} - \frac{D}{k_B T} \text{III}.
\end{aligned} \tag{135}$$

Using (19), for any smooth function $g(x, t)$, we have $\partial_t g + \nabla_x \cdot (ug) = D_t g$, where D_t is given by (23). It follows that $\text{I} = D_t(\rho_f Q_f)$ and, using (86), that

$$\text{I} = \rho_f D_t Q_f. \tag{136}$$

Using Stokes theorem, we get:

$$\begin{aligned}
\text{III}_{ij} &= \int_{\mathbb{S}^{n-1}} \nabla_\omega \cdot (f \nabla_\omega \mu_f^R) \omega_i \omega_j \, d\omega = - \int_{\mathbb{S}^{n-1}} f \nabla_\omega \mu_f^R \cdot \nabla_\omega (\omega_i \omega_j) \, d\omega \\
&= - \int_{\mathbb{S}^{n-1}} f \nabla_\omega \mu_f^R \cdot (\omega_j P_{\omega^\perp} e_i + \omega_i P_{\omega^\perp} e_j) \, d\omega \\
&= - \int_{\mathbb{S}^{n-1}} f (\omega_j \nabla_\omega \mu_f^R \cdot e_i + \omega_i \nabla_\omega \mu_f^R \cdot e_j) \, d\omega = - \frac{2}{\Lambda} ((\sigma_f^R)_s)_{ij},
\end{aligned} \tag{137}$$

where again, e_i denotes the i -th vector of the canonical basis of \mathbb{R}^n . Now, similarly to III, we have,

$$\begin{aligned}
\text{II}_{ij} &= \int_{\mathbb{S}^{n-1}} \nabla_\omega \cdot (f(\Lambda P_{\omega^\perp} E - W)\omega) \omega_i \omega_j \, d\omega \\
&= - \int_{\mathbb{S}^{n-1}} f ((\Lambda P_{\omega^\perp} E - W)\omega) \cdot (\omega_i e_j + \omega_j e_i) \, d\omega \\
&= - \int_{\mathbb{S}^{n-1}} f ((\Lambda E - W)(\omega \otimes \omega) + (\omega \otimes \omega)(\Lambda E + W) - 2\Lambda \omega^{\otimes 4} : E)_{ij} \, d\omega,
\end{aligned}$$

which leads to

$$\text{II} = \rho_f \left(-\Lambda(EQ_f + Q_f E) + WQ_f - Q_f W - \frac{2\Lambda}{n}E + 2\Lambda \mathbb{T}_f : E \right). \tag{138}$$

Finally, using (21), the antisymmetric part of σ_f^R is given by:

$$(\sigma_f^R)_a = \frac{1}{2} \int_{\mathbb{S}^{n-1}} (\omega \otimes \nabla_\omega U_f^R - \nabla_\omega U_f^R \otimes \omega) f \, d\omega. \tag{139}$$

Now, inserting (136), (137), (138) and (139) into (135) leads to (22). \square

Appendix B. Appendix to Section 3 on main result.

B.1. Proof of Prop. 2 on properties of S_2 . The proof uses Lemma 4.1 of [21] which we recall here without proof.

Lemma B.1. *Let $n \geq 2$. Define $\omega_\perp = P_{\Omega^\perp}\omega$. For any function $k: [-1, 1] \rightarrow \mathbb{R}$, $r \mapsto k(r)$, we have:*

$$\int_{\mathbb{S}^{n-1}} k(\omega \cdot \Omega) \omega_\perp^{\otimes(2k+1)} d\omega = 0, \quad \forall k \in \mathbb{N}, \tag{140}$$

$$\int_{\mathbb{S}^{n-1}} k(\omega \cdot \Omega) \omega_\perp \otimes \omega_\perp d\omega = \frac{1}{n-1} \int_{\mathbb{S}^{n-1}} k(\omega \cdot \Omega) (1 - (\omega \cdot \Omega)^2) d\omega P_{\Omega^\perp}. \tag{141}$$

Proof of Proposition 2. (i) The decomposition

$$\omega = (\omega \cdot \Omega) \Omega + \omega_\perp, \tag{142}$$

leads to

$$\omega \otimes \omega = (\omega \cdot \Omega)^2 \Omega \otimes \Omega + (\omega \cdot \Omega) (\omega_\perp \otimes \Omega + \Omega \otimes \omega_\perp) + \omega_\perp \otimes \omega_\perp. \tag{143}$$

We insert (143) into (10) with $f = \rho G_{\eta A_\Omega}$. Thanks to (45), $\rho G_{\eta A_\Omega}$ is a function of $\omega \cdot \Omega$ only. So, the contribution of the middle term of (143) vanishes thanks to (140) and the contribution of the last term can be computed using (141). Using that $P_{\Omega^\perp} = \text{Id} - \Omega \otimes \Omega$, we get

$$Q_{G_{\eta A_\Omega}} = \langle (\omega \cdot \Omega)^2 \rangle_{G_{\eta A_\Omega}} \Omega \otimes \Omega + \frac{1}{n-1} \langle 1 - (\omega \cdot \Omega)^2 \rangle_{G_{\eta A_\Omega}} (\text{Id} - \Omega \otimes \Omega) - \frac{1}{n} \text{Id}.$$

Rearranging these terms, we find (48).

(ii) The leading eigenvalue of $Q_{G_{\eta A_\Omega}}$ is $\frac{n-1}{n} S_2(\eta)$ and is associated with the eigenvector Ω . Thus, by virtue of (12), the order parameter $\chi_{\rho G_{\eta A_\Omega}}$ is equal to $S_2(\eta)$.

(iii) We first compute $S_2(0)$. When $\eta = 0$, we have $G_{\eta A_\Omega} = 1$. Thus, $S_2(0) = \langle (n(\omega \cdot \Omega)^2 - 1)/(n-1) \rangle_1 =: r/s$, where, using the spherical coordinates as in the proof of Proposition 1, the numerator r is given by

$$r = \int_0^\pi (n \cos^2 \theta - 1) \sin^{n-2} \theta d\theta = (n-1)W_{n-2} - nW_n.$$

Here, W_n is twice the Wallis integral $W_n = \int_0^\pi \sin^n \theta d\theta$. From the well-known recursion formula for the Wallis integral (which can be easily proved by integration by parts): $W_n = \frac{n-1}{n} W_{n-2}$, we get that $r = 0$ and thus, that $S_2(0) = 0$.

We now show that $S_2'(\eta) \geq 0$, for all $\eta \geq 0$, where the prime denotes the derivative with respect to η . We have $S_2(\eta) = 1 - \frac{n}{n-1} \langle 1 - (\omega \cdot \Omega)^2 \rangle_{G_{\eta A_\Omega}} =: 1 - \frac{n}{n-1} F(\eta)$. We show that $F' \leq 0$. Using again spherical coordinates, we have $F = I_n/I_{n-2}$ with $I_n(\eta) = \int_0^{\pi/2} \exp(\eta \cos^2 \theta) \sin^n \theta d\theta$ (by symmetry, we can reduce the interval of integration to $[0, \pi/2]$). Thus, $F' = (I_n' I_{n-2} - I_{n-2}' I_n)/I_{n-2}^2 =: A/I_{n-2}^2$. We check the sign of the numerator A . We have

$$A(\eta) = \int_{[0, \frac{\pi}{2}]^2} e^{\eta(\cos^2 \theta + \cos^2 \theta')} \sin^{n-2} \theta \sin^{n-2} \theta' \sin^2 \theta (\cos^2 \theta - \cos^2 \theta') d\theta d\theta' = \frac{1}{2} \int_{[0, \frac{\pi}{2}]^2} e^{\eta(\cos^2 \theta + \cos^2 \theta')} \sin^{n-2} \theta \sin^{n-2} \theta' (\sin^2 \theta - \sin^2 \theta') (\cos^2 \theta - \cos^2 \theta') d\theta d\theta',$$

where we pass from the first to the second line by exchanging θ and θ' . Since \sin is increasing and \cos is decreasing on $[0, \frac{\pi}{2}]$, we have $A \leq 0$.

Finally, when $\eta \rightarrow \infty$, the measure $G_{\eta A \Omega} d\omega$ concentrates onto the sum of Dirac deltas $\frac{1}{2}(\delta_{\Omega} + \delta_{-\Omega})$. Since $P_2(\pm 1) = 1$, it follows that $S_2 \rightarrow 1$ when $\eta \rightarrow \infty$. This ends the proof. \square

B.2. Proof of Eq. (66) for the Leslie stresses. We have $f^\varepsilon \rightarrow f$ as $\varepsilon \rightarrow 0$ with f given by (60). We will abbreviate $G_{\eta A \Omega}$ into G for simplicity. We define

$$\sigma = \lim_{\varepsilon \rightarrow 0} (\zeta \rho_{f^\varepsilon} \mathbb{T}_{f^\varepsilon} : E^\varepsilon + \sigma_{f^\varepsilon}^1) = \zeta \rho \mathbb{T}_G : E + \sigma_{\rho G}^1.$$

From (32), we get

$$\begin{aligned} \sigma = \rho \left\{ \frac{\Lambda^2}{2} (EQ_G + Q_G E) + \frac{\Lambda}{2} (Q_G W - W Q_G) + \frac{\Lambda^2}{n} E \right. \\ \left. + (\zeta - \Lambda^2) \mathbb{T}_G : E - \frac{\Lambda}{2} D_t Q_G + \alpha \beta [\Delta_x (\rho Q_G) Q_G - Q_G \Delta_x (\rho Q_G)] \right\}. \end{aligned} \quad (144)$$

Now, for a generic distribution function f , we introduce the fourth-order tensorial order parameter given by

$$\mathbb{Q}_f = \mathbb{T}_f - \frac{6}{n+4} (\langle \omega \otimes \omega \rangle_f \otimes \text{Id})_s + \frac{3}{(n+2)(n+4)} (\text{Id} \otimes \text{Id})_s. \quad (145)$$

Here, $(\langle \omega \otimes \omega \rangle_f \otimes \text{Id})_s$ and $(\text{Id} \otimes \text{Id})_s$ denote the symmetrizations of the fourth-order tensors $\langle \omega \otimes \omega \rangle_f \otimes \text{Id}$ and $\text{Id} \otimes \text{Id}$ respectively. Specifically,

$$\begin{aligned} 6((\langle \omega \otimes \omega \rangle_f \otimes \text{Id} \otimes \text{Id})_s)_{ijkl} &= \langle \omega_i \omega_j \rangle_f \delta_{kl} + \langle \omega_i \omega_k \rangle_f \delta_{jl} + \langle \omega_i \omega_l \rangle_f \delta_{jk} \\ &\quad + \langle \omega_j \omega_k \rangle_f \delta_{il} + \langle \omega_j \omega_l \rangle_f \delta_{ik} + \langle \omega_k \omega_l \rangle_f \delta_{ij}, \\ 3((\text{Id} \otimes \text{Id})_s)_{ijkl} &= \delta_{ij} \delta_{kl} + \delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}, \end{aligned}$$

where δ denotes the Kronecker symbol. Eq. (145) corresponds to the decomposition of \mathbb{T}_f into irreducible tensors, i.e. invariant tensors under the action of the orthogonal group. The coefficients of the decomposition can be obtained by the requirement that the contraction of \mathbb{Q}_f with respect to any two indices is zero. Owing to the fact that $\langle \omega \otimes \omega \rangle_f = Q_f + \frac{1}{n} \text{Id}$, we get

$$\mathbb{Q}_f = \mathbb{T}_f - \frac{6}{n+4} (Q_f \otimes \text{Id})_s - \frac{3}{n(n+2)} (\text{Id} \otimes \text{Id})_s, \quad (146)$$

where the definition of $(Q_f \otimes \text{Id})_s$ is similar to that of $(\langle \omega \otimes \omega \rangle_f \otimes \text{Id} \otimes \text{Id})_s$. Then, using (146), we have

$$\mathbb{T}_f : E = \mathbb{Q}_f : E + \frac{2}{n+4} (EQ_f + Q_f E) + \frac{2}{n(n+2)} E + \frac{1}{n+4} (Q_f : E) \text{Id}.$$

Inserting this identity (with $f = G$) into (144), we get

$$\begin{aligned} \sigma = \rho \left\{ \left(\frac{\Lambda^2}{2} + \frac{2(\zeta - \Lambda^2)}{n+4} \right) (EQ_G + Q_G E) + \frac{\Lambda}{2} (Q_G W - W Q_G) \right. \\ \left. + \frac{1}{n} \left(\Lambda^2 + \frac{2(\zeta - \Lambda^2)}{n+2} \right) E + (\zeta - \Lambda^2) \mathbb{Q}_G : E + \frac{\zeta - \Lambda^2}{n+4} (Q_G : E) \text{Id} \right. \\ \left. - \frac{\Lambda}{2} D_t Q_G + \alpha \beta [\Delta_x (\rho Q_G) Q_G - Q_G \Delta_x (\rho Q_G)] \right\}. \end{aligned} \quad (147)$$

Now, we state two lemmas whose proofs are deferred to the end of the present proof

Lemma B.2. *We have*

$$\mathbb{Q}_{G_{\eta A_\Omega}} = S_4(\eta) \mathbb{A}_\Omega, \quad (148)$$

where $S_4(\eta)$ is given by (46) and where

$$\mathbb{A}_\Omega = \Omega^{\otimes 4} - \frac{6}{n+4}(\Omega \otimes \Omega \otimes \text{Id})_s + \frac{3}{(n+2)(n+4)}(\text{Id} \otimes \text{Id})_s. \quad (149)$$

Lemma B.3. *We have*

$$\begin{aligned} \rho \alpha \beta [\Delta_x(\rho Q_G) Q_G - Q_G \Delta_x(\rho Q_G)] &= \rho \frac{\Lambda S_2(\eta)}{2} \left[\frac{1}{c} (N \otimes \Omega - \Omega \otimes N) \right. \\ &\quad \left. - (E(\Omega \otimes \Omega) - (\Omega \otimes \Omega)E) \right], \end{aligned} \quad (150)$$

with N given by (68).

From (148) and (149), it follows that

$$\begin{aligned} \mathbb{Q}_G : E &= S_4 \left\{ (E : (\Omega \otimes \Omega)) \Omega \otimes \Omega - \frac{2}{n+4} [(\Omega \otimes \Omega)E + E(\Omega \otimes \Omega)] \right. \\ &\quad \left. - \frac{1}{n+4} (E : (\Omega \otimes \Omega)) \text{Id} + \frac{2}{(n+2)(n+4)} E \right\}, \end{aligned} \quad (151)$$

where the dependence of S_4 on η is omitted for simplicity. Likewise, with (48), we get

$$EQ_G + Q_GE = S_2 [(\Omega \otimes \Omega)E + E(\Omega \otimes \Omega) - \frac{2}{n}E], \quad (152)$$

$$Q_G W - W Q_G = S_2 [(\Omega \otimes \Omega)W - W(\Omega \otimes \Omega)], \quad (153)$$

$$D_t Q_G = S_2 [D_t \Omega \otimes \Omega + \Omega \otimes D_t \Omega]. \quad (154)$$

In (154), we have used that $D_t S_2(\eta(\rho)) = \frac{dS_2}{d\eta}(\eta(\rho)) \frac{d\eta}{d\rho}(\rho) D_t \rho = 0$ thanks to (86). Inserting Eqs. (150) to (154) into (147), we get $\sigma = \sigma_L + \nabla_x \varphi$ where φ is a scalar function which can be absorbed in the pressure, and σ_L is given by (66) with the constants, α_k , $k = 1, \dots, 6$ given by (55)-(58). This ends the proof. \square

Proof of Lemma B.2. Using (146), (48), (46) and (47), we get that

$$\begin{aligned} \mathbb{Q}_{G_{\eta A_\Omega}} &= \mathbb{T}_{G_{\eta A_\Omega}} - \frac{6S_2}{n+4} (\Omega \otimes \Omega \otimes \text{Id})_s + \left(\frac{6S_2}{n(n+4)} - \frac{3}{n(n+2)} \right) (\text{Id} \otimes \text{Id})_s \\ &= \mathbb{T}_{G_{\eta A_\Omega}} - \frac{6(n\langle X^2 \rangle - 1)}{(n-1)(n+4)} (\Omega \otimes \Omega \otimes \text{Id})_s \\ &\quad + \left(\frac{6(n\langle X^2 \rangle - 1)}{(n-1)n(n+4)} - \frac{3}{n(n+2)} \right) (\text{Id} \otimes \text{Id})_s, \end{aligned} \quad (155)$$

where $X = \omega \cdot \Omega$ and where we drop the index $G_{\eta A_\Omega}$ on the brackets $\langle \cdot \rangle$. Now, using the decomposition (142), we get

$$\mathbb{T}_{G_{\eta A_\Omega}} = \langle X^4 \rangle \Omega^{\otimes 4} + \left(\langle X^2 \omega_\perp \otimes \omega_\perp \rangle \otimes (\Omega \otimes \Omega) \right)_s + \langle \omega_\perp^{\otimes 4} \rangle.$$

We use (141) to compute $\langle X^2 \omega_\perp \otimes \omega_\perp \rangle$. To evaluate $\langle \omega_\perp^{\otimes 4} \rangle$ we recall the last part of Lemma 4.1 of [21] without proof: with the notations of Lemma B.1, we have

$$\int_{\mathbb{S}^{n-1}} k(\omega \cdot \Omega) \omega_\perp^{\otimes 4} d\omega = \int_{\mathbb{S}^{n-1}} \frac{3k(\omega \cdot \Omega) (1 - (\omega \cdot \Omega)^2)^2}{(n-1)(n+1)} d\omega (P_{\Omega^\perp} \otimes P_{\Omega^\perp})_s.$$

This leads to

$$\mathbb{T}_{G_{\eta A_\Omega}} = \langle X^4 \rangle \Omega^{\otimes 4} + \frac{6 \langle X^2(1-X^2) \rangle}{n-1} (\Omega \otimes \Omega \otimes P_{\Omega^\perp})_s + \frac{3 \langle (1-X^2)^2 \rangle}{(n-1)(n+1)} (P_{\Omega^\perp} \otimes P_{\Omega^\perp})_s.$$

Using that $P_{\Omega^\perp} = \text{Id} - \Omega \otimes \Omega$, we obtain

$$\begin{aligned} \mathbb{T}_{G_{\eta A_\Omega}} &= \left(\langle X^4 \rangle - \frac{6 \langle X^2(1-X^2) \rangle}{n-1} + \frac{3 \langle (1-X^2)^2 \rangle}{(n-1)(n+1)} \right) \Omega^{\otimes 4} \\ &\quad + \left(\frac{6 \langle X^2(1-X^2) \rangle}{n-1} - \frac{6 \langle (1-X^2)^2 \rangle}{(n-1)(n+1)} \right) (\Omega \otimes \Omega \otimes \text{Id})_s \\ &\quad + \frac{3 \langle (1-X^2)^2 \rangle}{(n-1)(n+1)} (\text{Id} \otimes \text{Id})_s. \end{aligned} \quad (156)$$

Now, inserting (156) into (155), we get (148). \square

Proof of Lemma B.3. Thanks to (50) and (129), we have

$$\begin{aligned} \alpha^2 \rho \Delta_x(\rho Q_G) Q_G &= \eta \Delta_x(\eta A_\Omega) A_\Omega \\ &= \eta \left[\Delta_x \eta A_\Omega^2 + \frac{2(n-1)}{n} (\nabla_x \eta \cdot \nabla_x) \Omega \otimes \Omega - \frac{2}{n} \Omega \otimes (\nabla_x \eta \cdot \nabla_x) \Omega - \frac{2\eta}{n} \nabla_x \Omega^T \nabla_x \Omega \right. \\ &\quad \left. + \frac{\eta(n-1)}{n} \Delta_x \Omega \otimes \Omega - \frac{\eta}{n} \Omega \otimes \Delta_x \Omega + \eta(\Omega \cdot \Delta_x \Omega) \Omega \otimes \Omega \right]. \end{aligned} \quad (157)$$

Let M be the tensor given by the left-hand side of (150). Using (62) and (49), it follows from (157) that

$$\begin{aligned} M &= \frac{\beta\eta}{\alpha} \left[2((\nabla_x \eta \cdot \nabla_x) \Omega \otimes \Omega - \Omega \otimes (\nabla_x \eta \cdot \nabla_x) \Omega) + \eta(\Delta_x \Omega \otimes \Omega - \Omega \otimes \Delta_x \Omega) \right] \\ &= \frac{\beta}{\alpha} \left[\Delta_x(\eta \Omega) \otimes (\eta \Omega) - (\eta \Omega) \otimes \Delta_x(\eta \Omega) \right] \\ &= \frac{\Lambda}{2\alpha} \left[\left(\frac{N}{c} - P_{\Omega^\perp} E \Omega \right) \otimes (\eta \Omega) - (\eta \Omega) \otimes \left(\frac{N}{c} - P_{\Omega^\perp} E \Omega \right) \right] \\ &= \frac{\rho \Lambda S_2}{2} \left[\left(\frac{N}{c} - P_{\Omega^\perp} E \Omega \right) \otimes \Omega - \Omega \otimes \left(\frac{N}{c} - P_{\Omega^\perp} E \Omega \right) \right]. \end{aligned} \quad (158)$$

Then, we note that there exists a real number z such that

$$(P_{\Omega^\perp} E \Omega) \otimes \Omega = (E \Omega) \otimes \Omega + z \Omega \otimes \Omega = E(\Omega \otimes \Omega) + z \Omega \otimes \Omega,$$

and that the same real number z is involved in the expression of $\Omega \otimes (P_{\Omega^\perp} E \Omega)$, so that we get

$$(P_{\Omega^\perp} E \Omega) \otimes \Omega - \Omega \otimes (P_{\Omega^\perp} E \Omega) = E(\Omega \otimes \Omega) - (\Omega \otimes \Omega) E.$$

Inserting this expression into (158), we get (150) which ends the proof of the Lemma. \square

B.3. Proof of Eq. (67) for the Ericksen stresses. We now compute $\lim_{\varepsilon \rightarrow 0} F_{f^\varepsilon}^1 = F_{\rho G_{\eta A_\Omega}}^1$. Thanks to (33), (30) and (51), we have

$$\begin{aligned} F_{\rho G_{\eta A_\Omega}}^1 &= -\rho \langle \nabla_x \mu_{\rho G_{\eta A_\Omega}}^1 \rangle_{G_{\eta A_\Omega}} \\ &= \beta \rho \left\{ \langle \nabla_x \Delta_x(\eta(\omega \cdot \Omega)^2) \rangle_{G_{\eta A_\Omega}} - \frac{1}{n} \nabla_x \Delta_x[\eta + (n-1)\alpha\rho] \right\}. \end{aligned} \quad (159)$$

We compute, using the repeated index summation convention:

$$\begin{aligned}
\partial_{x_i} [\Delta_x (\eta (\omega \cdot \Omega)^2)] &= 2(\omega \cdot \Omega) \partial_{x_i} \Omega_k \omega_{\perp k} \partial_{x_j x_j}^2 \eta + (\omega \cdot \Omega)^2 \partial_{x_i x_j x_j}^3 \eta \\
&+ 4 \partial_{x_i} \Omega_k \omega_{\perp k} \partial_{x_j} \Omega_\ell \omega_{\perp \ell} \partial_{x_j} \eta + 4(\omega \cdot \Omega) \partial_{x_i x_j}^2 \Omega_k \omega_k \partial_{x_j} \eta \\
&+ 4(\omega \cdot \Omega) \partial_{x_j} \Omega_k \omega_{\perp k} \partial_{x_i x_j}^2 \eta + 2 \partial_{x_j} \Omega_k \omega_{\perp k} \partial_{x_j} \Omega_\ell \omega_{\perp \ell} \partial_{x_i} \eta \\
&+ 4 \partial_{x_i x_j}^2 \Omega_k \omega_k \partial_{x_j} \Omega_\ell \omega_{\perp \ell} \eta + 2(\omega \cdot \Omega) \partial_{x_j x_j}^2 \Omega_k \omega_k \partial_{x_i} \eta \\
&+ 2 \partial_{x_j x_j}^2 \Omega_k \omega_k \partial_{x_i} \Omega_\ell \omega_{\perp \ell} \eta + 2(\omega \cdot \Omega) \partial_{x_i x_j x_j}^3 \Omega_k \omega_k \eta. \quad (160)
\end{aligned}$$

Thanks to (46) and (47), we have the following identities

$$\langle (\omega \cdot \Omega)^2 \rangle_{G_{\eta A \Omega}} = \frac{(n-1)S_2 + 1}{n}, \quad \langle 1 - (\omega \cdot \Omega)^2 \rangle_{G_{\eta A \Omega}} = \frac{n-1}{n}(1 - S_2).$$

Furthermore, the decomposition (142) and the fact that $|\Omega|^2 = 1$, lead to the following identities

$$\begin{aligned}
\partial_{x_i} \Omega_k \Omega_k &= 0, \quad \partial_{x_i x_j}^2 \Omega_k \Omega_k = -\partial_{x_i} \Omega_k \partial_{x_j} \Omega_k, \\
(P_{\omega^\perp})_{k\ell} \partial_{x_j} \Omega_\ell &= \partial_{x_j} \Omega_k, \quad \partial_{x_i x_j}^2 \Omega_k \partial_{x_j} \Omega_k = \frac{1}{2} \partial_{x_i} (\partial_{x_j} \Omega_k \partial_{x_j} \Omega_k) \\
\partial_{x_i x_j x_j}^3 \Omega_k \Omega_k &= -\partial_{x_j} (\partial_{x_i} \Omega_k \partial_{x_j} \Omega_k) - \frac{1}{2} \partial_{x_i} (\partial_{x_j} \Omega_k \partial_{x_j} \Omega_k), \\
\partial_{x_j x_j}^2 \Omega_k \partial_{x_i} \Omega_k &= \partial_{x_j} (\partial_{x_i} \Omega_k \partial_{x_j} \Omega_k) - \frac{1}{2} \partial_{x_i} (\partial_{x_j} \Omega_k \partial_{x_j} \Omega_k).
\end{aligned}$$

Thus, taking the bracket $\langle \cdot \rangle_{G_{\eta A \Omega}}$ of (160), noting that all odd powers of $\omega \cdot \Omega$ or of ω_{\perp} vanish by antisymmetry and using (141) and the previous identities, we finally get

$$\begin{aligned}
\langle \nabla_x \Delta_x (\eta (\omega \cdot \Omega)^2) \rangle_{G_{\eta A \Omega}} &= \frac{(n-1)S_2 + 1}{n} \nabla_x \Delta_x \eta \\
&- 2S_2 (2\nabla_x \Omega \nabla_x \Omega^T + |\nabla_x \Omega|^2 \text{Id}) \nabla_x \eta \\
&- S_2 \nabla_x \cdot (2\nabla_x \Omega \nabla_x \Omega^T + |\nabla_x \Omega|^2 \text{Id}) \eta.
\end{aligned}$$

Inserting this equation into (159), using (49) and noting that for a $n \times n$ tensor S and a scalar φ , we have $\nabla_x \cdot (S\varphi) = (\nabla_x \cdot S)\varphi + S^T \nabla_x \varphi$, we get

$$\begin{aligned}
F_{\rho G_{\eta A \Omega}}^1 &= -\frac{\beta}{\alpha} \nabla_x \cdot \left[\eta^2 (2\nabla_x \Omega \nabla_x \Omega^T + |\nabla_x \Omega|^2 \text{Id}) \right] \\
&+ \beta \left\{ \frac{n-1}{n\alpha} \eta \nabla_x \Delta_x \eta - \frac{(n-1)\alpha}{n} \rho \nabla_x \Delta_x \rho \right\}. \quad (161)
\end{aligned}$$

The first of the following identities follows again from the fact that $|\Omega|^2 = 1$ and the second one is just straightforward algebra (which will also be applied with ρ replacing η):

$$\begin{aligned}
\eta^2 \nabla_x \Omega \nabla_x \Omega^T &= \nabla_x (\eta \Omega) \nabla_x (\eta \Omega)^T - \nabla_x \eta \otimes \nabla_x \eta, \\
\eta \nabla_x \Delta_x \eta &= -\nabla_x \cdot (\nabla_x \eta \otimes \nabla_x \eta) + \nabla_x (\eta \Delta_x \eta + \frac{1}{2} |\nabla_x \eta|^2).
\end{aligned}$$

Inserting these identities into (161), we get $F_{\rho G_{\eta A \Omega}}^1 = \nabla_x \cdot \sigma_E + \nabla_x \varphi$, where σ_E is given by (67) and φ is a scalar function (different from the one appearing at the end of Section B.2) which can be absorbed in the pressure p . This ends the proof. \square

B.4. Proof of the energy identity (74). Taking the dot product of (63) with u , integrating with respect to x on \mathbb{R}^n and using Stokes formula assuming that the spatial boundary terms vanish at infinity, we get

$$\frac{d}{dt} \int_{\mathbb{R}^n} \frac{|u|^2}{2} dx + \frac{1}{\text{Re}} \int_{\mathbb{R}^n} |\nabla_x u|^2 dx + \frac{1}{\text{Re} E \Gamma} \int_{\mathbb{R}^n} (\sigma_L + \sigma_E) : \nabla_x u dx = 0. \quad (162)$$

We first compute the contribution of the Leslie stresses. Using the symmetry of E and $\Omega \otimes \Omega$, we first have

$$[\alpha_1(E : (\Omega \otimes \Omega))\Omega \otimes \Omega + \alpha_4 E] : \nabla_x u = \alpha_1(E : (\Omega \otimes \Omega))^2 + \alpha_4 |E|^2. \quad (163)$$

Then, we remark that

$$\begin{aligned} ((\Omega \otimes \Omega)E) : \nabla_x u &= |E\Omega|^2 - (E\Omega) \cdot (W\Omega), \\ (E(\Omega \otimes \Omega)) : \nabla_x u &= |E\Omega|^2 + (E\Omega) \cdot (W\Omega), \end{aligned}$$

which, with the second equation (69), gives

$$[\alpha_5(\Omega \otimes \Omega)E + \alpha_6 E(\Omega \otimes \Omega)] : \nabla_x u = (\alpha_5 + \alpha_6)|E\Omega|^2 + \gamma_2(E\Omega) \cdot (W\Omega). \quad (164)$$

Also, with (70), we have

$$N = -\frac{\gamma_2}{\gamma_1} P_{\Omega^\perp} E\Omega + \frac{1}{\gamma_1} P_{\Omega^\perp} H := N_1 + N_2.$$

Remarking that $(\Omega \otimes \Omega) : W = 0$ by the antisymmetry of W , we get

$$\begin{aligned} (\Omega \otimes N_1) : \nabla_x u &= -\frac{\gamma_2}{\gamma_1} [\Omega \otimes (E\Omega - (\Omega \cdot E\Omega)\Omega)] : (E + W) \\ &= -\frac{\gamma_2}{\gamma_1} [|E\Omega|^2 - (E : (\Omega \otimes \Omega))^2 - (E\Omega) \cdot (W\Omega)], \end{aligned}$$

and similarly

$$(N_1 \otimes \Omega) : \nabla_x u = -\frac{\gamma_2}{\gamma_1} [|E\Omega|^2 - (E : (\Omega \otimes \Omega))^2 + (E\Omega) \cdot (W\Omega)],$$

which, using (69), gives

$$[\alpha_2 \Omega \otimes N_1 + \alpha_3 N_1 \otimes \Omega] : \nabla_x u = -\frac{\gamma_2^2}{\gamma_1} [|E\Omega|^2 - (E : (\Omega \otimes \Omega))^2] - \gamma_2(E\Omega) \cdot (W\Omega). \quad (165)$$

Then, using (69), we compute

$$\begin{aligned} [\alpha_2 \Omega \otimes N_2 + \alpha_3 N_2 \otimes \Omega] : \nabla_x u &= \frac{1}{\gamma_1} [\alpha_2 \Omega \otimes P_{\Omega^\perp} H + \alpha_3 P_{\Omega^\perp} H \otimes \Omega] : (E + W) \\ &= \frac{1}{\gamma_1} [(\alpha_2 + \alpha_3)(P_{\Omega^\perp} H \otimes \Omega) : E + (\alpha_3 - \alpha_2)(P_{\Omega^\perp} H \otimes \Omega) : W] \\ &= P_{\Omega^\perp} H \cdot \left[\frac{\gamma_2}{\gamma_1} E\Omega + W\Omega \right] = P_{\Omega^\perp} H \cdot P_{\Omega^\perp} \left[\frac{\gamma_2}{\gamma_1} E\Omega + W\Omega \right] \\ &= \frac{1}{\gamma_1} |P_{\Omega^\perp} H|^2 - H \cdot (\partial_t \Omega + u \cdot \nabla_x \Omega), \end{aligned} \quad (166)$$

where, for the last equality, we have used (70) and (68) and the fact that $\partial_t \Omega + u \cdot \nabla_x \Omega$ is normal to Ω . Then, collecting (163) to (166) and using (66) leads to

$$\begin{aligned} \int_{\mathbb{R}^n} \sigma_L : \nabla_x u dx &= \int_{\mathbb{R}^n} \rho \left\{ \left(\alpha_1 + \frac{\gamma_2^2}{\gamma_1} \right) (E : (\Omega \otimes \Omega))^2 + \alpha_4 |E|^2 \right. \\ &\quad \left. + \left(\alpha_5 + \alpha_6 - \frac{\gamma_2^2}{\gamma_1} \right) |E\Omega|^2 + \frac{1}{\gamma_1} |P_{\Omega^\perp} H|^2 - H \cdot (\partial_t \Omega + u \cdot \nabla_x \Omega) \right\} dx. \end{aligned} \quad (167)$$

Expression (67) for the Ericksen stresses involves three terms which we will denote by σ_E^Ω , σ_E^η , σ_E^ρ in the order in which they appear in this expression. We compute the contribution of each term successively. We have, using Stokes's formula, (73), (64) and assuming that the boundary terms vanish at infinity:

$$\begin{aligned}
\int_{\mathbb{R}^n} \sigma_E^\Omega : \nabla_x u \, dx &= -\frac{2\beta}{\alpha} \int_{\mathbb{R}^n} (\nabla_x(\eta\Omega)\nabla_x(\eta\Omega)^T) : \nabla_x u \, dx \\
&= \frac{2\beta}{\alpha} \int_{\mathbb{R}^n} \left\{ \Delta_x(\eta\Omega) \cdot ((u \cdot \nabla_x)(\eta\Omega)) + \nabla_x(\eta\Omega) \cdot ((u \cdot \nabla_x)(\nabla_x(\eta\Omega))^T) \right\} dx \\
&= \frac{2\beta}{\alpha} \int_{\mathbb{R}^n} \left\{ \eta \Delta_x(\eta\Omega) \cdot ((u \cdot \nabla_x)\Omega) + \Delta_x(\eta\Omega) \cdot \Omega (u \cdot \nabla_x)\eta + \nabla_x \cdot \left(u \frac{|\nabla_x(\eta\Omega)|^2}{2} \right) \right\} dx \\
&= \int_{\mathbb{R}^n} \rho H \cdot ((u \cdot \nabla_x)\Omega) \, dx + \frac{2\beta}{\alpha} \int_{\mathbb{R}^n} \Delta_x(\eta\Omega) \cdot \Omega (u \cdot \nabla_x)\eta \, dx. \tag{168}
\end{aligned}$$

A similar computation gives

$$\int_{\mathbb{R}^n} \sigma_E^\eta : \nabla_x u \, dx = -\frac{(n+1)\beta}{n\alpha} \int_{\mathbb{R}^n} \Delta_x \eta (u \cdot \nabla_x)\eta \, dx, \tag{169}$$

$$\int_{\mathbb{R}^n} \sigma_E^\rho : \nabla_x u \, dx = -\frac{(n-1)\alpha\beta}{n} \int_{\mathbb{R}^n} \Delta_x \rho (u \cdot \nabla_x)\rho \, dx. \tag{170}$$

Now, we consider the Oseen-Frank energy and successively compute the time derivative of each of the terms in (72). We first have, thanks to Stokes's formula:

$$\frac{d\mathcal{E}_F^\Omega}{dt} = \frac{2\beta}{\alpha} \int_{\mathbb{R}^n} \text{Tr} \left\{ \nabla_x(\eta\Omega) (\partial_t \nabla_x(\eta\Omega))^T \right\} dx = -\frac{2\beta}{\alpha} \int_{\mathbb{R}^n} \Delta_x(\eta\Omega) \cdot \partial_t(\eta\Omega) \, dx,$$

With (73), this leads to:

$$\frac{d\mathcal{E}_F^\Omega}{dt} + \int_{\mathbb{R}^n} \rho H \cdot \partial_t \Omega \, dx + \frac{2\beta}{\alpha} \int_{\mathbb{R}^n} \Delta_x(\eta\Omega) \cdot \Omega \partial_t \eta \, dx = 0. \tag{171}$$

Straightforwardly, we get

$$\frac{d\mathcal{E}_F^\eta}{dt} - \frac{(n+1)\beta}{n\alpha} \int_{\mathbb{R}^n} \Delta_x \eta \cdot \partial_t \eta \, dx = 0, \tag{172}$$

$$\frac{d\mathcal{E}_F^\rho}{dt} - \frac{(n-1)\alpha\beta}{n} \int_{\mathbb{R}^n} \Delta_x \rho \cdot \partial_t \rho \, dx = 0. \tag{173}$$

Now, adding (162), (171), (172), (173) together, using (167), (168), (169), (170) to eliminate σ_L and σ_E and finally using that $D_t \rho = 0$ and $D_t \eta = \frac{d\eta}{d\rho} D_t \rho = 0$, we get Eq. (74). \square

Appendix C. Appendix to Section 5 on GCI.

C.1. Proof of Proposition 10. We first note that Eq. (97) which defines h_η can be alternately written as (dropping the index η for simplicity):

$$(1-r^2)h'' + (2\eta(1-r^2) - (n+1))rh' - (2\eta r^2 + n-1)h = r. \tag{174}$$

With (98), we have

$$h(r) = -\frac{1}{2\eta} \frac{1}{\sqrt{1-r^2}} g(\cos^{-1} r).$$

Then,

$$\begin{aligned} h'(r) &= -\frac{1}{2\eta} \left[-\frac{g'(\cos^{-1} r)}{1-r^2} + \frac{r g(\cos^{-1} r)}{(1-r^2)^{3/2}} \right], \\ h''(r) &= -\frac{1}{2\eta} \left[\frac{g''(\cos^{-1} r)}{(1-r^2)^{3/2}} - \frac{3r g'(\cos^{-1} r)}{(1-r^2)^2} + \frac{(1+2r^2) g(\cos^{-1} r)}{(1-r^2)^{5/2}} \right]. \end{aligned}$$

Inserting these expressions in (174) and changing r into $\cos \theta$, we get

$$g'' + \frac{\cos \theta (n-2-2\eta \sin^2 \theta)}{\sin \theta} g' - \frac{n-2}{\sin^2 \theta} g = -2\eta \cos \theta \sin \theta, \quad (175)$$

But, we have

$$\frac{1}{\sin^{n-2} \theta} (\sin^{n-2} \theta g')' = g'' + (n-2) \frac{\cos \theta}{\sin \theta} g'.$$

With this and (52), we realize that (175) is nothing but (53). \square

C.2. Proof of Lemma 5.10. We use the same notations as in the proof of Lemma B.2. From (156), we get

$$\begin{cases} a_1 = \langle X^4 \rangle - \frac{6 \langle X^2(1-X^2) \rangle}{n-1} + \frac{3 \langle (1-X^2)^2 \rangle}{(n-1)(n+1)}, & (176) \\ a_2 = \frac{\langle X^2(1-X^2) \rangle}{n-1} - \frac{\langle (1-X^2)^2 \rangle}{(n-1)(n+1)}, & (177) \\ a_3 = \frac{\langle (1-X^2)^2 \rangle}{(n-1)(n+1)}, \end{cases}$$

Thus, with the change to spherical coordinates used in the proof of Prop. 1, an integration by parts, and Eqs. (46), (47) and (49), we get

$$\begin{aligned} a_2 + a_3 &= \frac{\langle X^2(1-X^2) \rangle}{n-1} = \frac{C_n}{(n-1)Z_\eta} \int_0^\pi e^{n \cos^2 \theta} \cos^2 \theta \sin^n \theta d\theta \\ &= \frac{C_n}{2(n-1)\eta Z_\eta} \int_0^\pi e^{n \cos^2 \theta} (n \cos^2 \theta - 1) \sin^{n-2} \theta d\theta \\ &= \frac{1}{2\eta} \frac{\langle nX^2 - 1 \rangle}{n-1} = \frac{S_2(\eta)}{2\eta} = \frac{1}{2\alpha\rho}, \end{aligned} \quad (178)$$

which shows the equality in (117).

Now, we have, thanks to (178)

$$(n-1)(n+1)a_2 = (n+2)\langle X^2(1-X^2) \rangle - \langle 1-X^2 \rangle = \frac{n+2}{2\eta} \langle nX^2 - 1 \rangle - \langle 1-X^2 \rangle.$$

Thanks to (46), (47), we have $\langle X^2 \rangle = \frac{1}{n}(1 + (n-1)S_2(\eta))$. So,

$$\begin{aligned} 2n(n+1)\eta a_2 &= \frac{n}{n-1} [(n(n+2) + 2\eta)\langle X^2 \rangle - (n+2 + 2\eta)] \\ &= (n(n+2) + 2\eta)S_2(\eta) - 2\eta. \end{aligned}$$

Thus, with the change to spherical coordinates used in the proof of Prop. 1, we have

$$\begin{aligned} 2n(n-1)(n+1)\eta Z_\eta C_n^{-1} a_2 &= \\ &= \int_0^\pi e^{\eta \cos^2 \theta} \left[(n(n+2) + 2\eta)(n \cos^2 \theta - 1) - 2(n-1)\eta \right] \sin^{n-2} \theta d\theta \\ &= \int_0^\pi e^{\eta \cos^2 \theta} \left[n(n+2)(n \cos^2 \theta - 1) + 2n\eta(\cos^2 \theta - 1) \right] \sin^{n-2} \theta d\theta \\ &= \int_0^\pi e^{\eta \cos^2 \theta} \left[2n(n+2)\eta \cos^2 \theta \sin^2 \theta - 2n\eta \sin^2 \theta \right] \sin^{n-2} \theta d\theta \\ &= 2\eta n \int_0^\pi e^{\eta \cos^2 \theta} ((n+2) \cos^2 \theta - 1) \sin^n \theta d\theta \end{aligned}$$

The passage between the third and fourth lines uses the same integration by parts as in (178). The other equalities are just simple algebraic rearrangements. Comparing with (46), (47), we notice that the integral of the last line is equal to the quantity $S_2^{(n+2)}$ which is the quantity S_2 in dimension $n+2$ up to a prefactor $(n+1)Z_\eta C_n^{-1}$. Thus, we have $a_2 = S_2^{(n+2)}/(n-1)$. Now, we can apply Prop. 2 (iii) and conclude that $0 < a_2 < \frac{1}{n-1}$. In particular, $a_2 \neq 0$, which finishes to show (117).

Finally, it is a simple algebra, using (176) and (177) to show that

$$a_1 + (n+4)a_2 = \frac{\langle nX^2 - 1 \rangle}{n-1} = S_2(\eta),$$

showing (118). This ends the proof. □

C.3. Proof of Prop. 12. Let $f = \rho G_{\eta A_\Omega}$ with $\eta \neq \eta(\rho)$. This means that (49) is not satisfied. In other words,

$$\eta' = \alpha \rho S_2(\eta) \neq \eta. \tag{179}$$

From (48), it follows that $\alpha \rho Q_{\rho G_{\eta A_\Omega}} = \eta' A_\Omega$. So, with (104), we get

$$D_{\rho G_{\eta A_\Omega}} C^* g(\omega) = L_{\eta' A_\Omega}^* g(\omega) - \alpha \rho (\rho Q)_{G_{\eta A_\Omega}} L_{\eta A_\Omega}^* g : \omega \otimes \omega. \tag{180}$$

Suppose that g is a GCI associated with (η, A_Ω) . Then, by (90), there exists $V \in \{\Omega\}^\perp$ such that

$$G_{\eta A_\Omega} L_{\eta A_\Omega}^* g = (\omega \cdot \Omega) (\omega \cdot V) G_{\eta A_\Omega}. \tag{181}$$

By a similar computation (using the same notations) to what was done in the proof of Lemma 5.9, we get

$$\begin{aligned} \alpha \rho (\rho Q)_{G_{\eta A_\Omega}} L_{\eta A_\Omega}^* g : \omega \otimes \omega &= 2\alpha \rho (a_2 + a_3) (\omega \cdot \Omega) (\omega \cdot V) \\ &= \frac{\eta'}{\eta} (\omega \cdot \Omega) (\omega \cdot V). \end{aligned} \tag{182}$$

For the second equality, we have used that $a_2 + a_3 = \frac{S_2(\eta)}{\eta}$ (see the proof of Lemma 5.10 in Appendix C.2) and (179). On the other hand, simple algebraic manipulations and the use of (181) show that

$$\begin{aligned} L_{\eta' A_\Omega}^* g(\omega) &= L_{\eta A_\Omega}^* g(\omega) + 2(\eta' - \eta) (\omega \cdot \Omega) P_{\omega^\perp} \Omega \cdot \nabla_\omega g \\ &= (\omega \cdot \Omega) (\omega \cdot V) + 2(\eta' - \eta) (\omega \cdot \Omega) P_{\omega^\perp} \Omega \cdot \nabla_\omega g. \end{aligned} \tag{183}$$

Inserting (182) and (183) into (180) gives

$$D_{\rho G_{\eta A\Omega}} C^* g(\omega) = (\eta' - \eta)(\omega \cdot \Omega) \left[-\frac{1}{\eta}(\omega \cdot V) + 2P_{\omega^\perp} \Omega \cdot \nabla_{\omega} g \right].$$

Suppose now that g is also an element of $\ker(D_{\rho G_{\eta A\Omega}} C^*)$. This implies that

$$2P_{\omega^\perp} \Omega \cdot \nabla_{\omega} g = \frac{1}{2\eta}(\omega \cdot V). \quad (184)$$

From now on, we restrict to dimension $n = 3$ and use the spherical coordinates (θ, φ) associated to the cartesian basis (V, W, Ω) with pole at Ω (defining $W = \Omega \times V$, using the symbol \times for the cross product). In these coordinates, (184) is written in terms of $\tilde{g}(\theta, \varphi) = g(\omega)$ according to

$$\partial_{\theta} \tilde{g} = \frac{1}{2\eta} \cos \varphi.$$

Thus,

$$\tilde{g}(\theta, \varphi) = \frac{1}{2\eta} \theta \cos \varphi + h(\varphi),$$

where h is an arbitrary function. The smoothness of g at $\omega = \Omega$ requires $h = 0$. However, we see that g cannot be smooth at $\omega = -\Omega$ (i.e. for $\theta = \pi$) because the function $\theta \cos \varphi$ does not tend to a constant when $\theta \rightarrow \pi$. However, by the elliptic regularity theorem, $g \in C^\infty(\mathbb{S}^{n-1})$. This is a contradiction. This means that the only possible solution is when $V = 0$, i.e.

$$\mathcal{C}_{\eta A\Omega} \cap \ker(D_{\rho G_{\eta A\Omega}} C^*) = \{0\}.$$

Since $\mathcal{C}_{\eta A\Omega} \neq \{0\}$, this shows (103) (with $\Sigma = A\Omega$) and ends the proof. \square

Appendix D. Appendix to Section 6 on the derivation of the equation for Ω .

D.1. **Proof of Eq. (132).** We first consider $\tilde{\gamma}_3$. With (127), (128) and (125), and using the spherical coordinates and the notations C_n and Z_η described in the proof of Prop. 1 as well as the change $r = \cos \theta$, we have (dropping the indices $\eta A\Omega$ to G and η to h for simplicity):

$$\begin{aligned} \tilde{\gamma}_3 &= \frac{\rho}{n-1} \int_{\mathbb{S}^{n-1}} G h(\omega \cdot \Omega) (1 - (\omega \cdot \Omega)^2) (2\eta(1 - 2(\omega \cdot \Omega)^2) - 2n) d\omega \\ &= \frac{\rho}{n-1} \frac{C_n}{Z_\eta} \int_0^1 (1 - r^2)^{\frac{n-1}{2}} e^{\eta r^2} r (2\eta(1 - 2r^2) - 2n) h dr, \end{aligned} \quad (185)$$

Besides, multiplying Eq. (97) by r , integrating with respect to $r \in [0, 1]$, and noting that, thanks to two successive integration by parts we have

$$\int_0^1 r \left((1 - r^2)^{\frac{n+1}{2}} e^{\eta r^2} h' \right)' dr = \int_0^1 (1 - r^2)^{\frac{n-1}{2}} e^{\eta r^2} r (2\eta(1 - r^2) - (n+1)) h dr,$$

we get

$$\int_0^1 (1 - r^2)^{\frac{n-1}{2}} e^{\eta r^2} r (2\eta(1 - 2r^2) - 2n) h dr = \int_0^1 (1 - r^2)^{\frac{n-1}{2}} e^{\eta r^2} r^2 dr. \quad (186)$$

Inserting (186) into (185) and integrating by parts once more, we get

$$\begin{aligned}\tilde{\gamma}_3 &= \frac{\rho}{n-1} \frac{C_n}{Z_\eta} \int_0^1 (1-r^2)^{\frac{n-1}{2}} e^{\eta r^2} r^2 dr \\ &= \frac{\rho}{n-1} \frac{C_n}{Z_\eta} \frac{1}{2\eta} \int_0^1 (1-r^2)^{\frac{n-3}{2}} e^{\eta r^2} (nr^2 - 1) dr \\ &= \frac{\rho}{2\eta} \int_{S^{n-1}} G \frac{n(\omega \cdot \Omega)^2 - 1}{n-1} d\omega = \frac{\rho \langle P_2(\omega \cdot \Omega) \rangle_G}{2\eta} = \frac{\rho S_2(\eta)}{2\eta},\end{aligned}$$

where, in the last line, we have reverted back to the variable ω and used (46) and (47). This shows the first equation in Formula (132).

We now consider $\tilde{\gamma}_1$. Changing to spherical coordinates in (125), we get

$$\tilde{\gamma}_1 = \frac{2\eta\rho}{n-1} \frac{C_n}{Z_\eta} \int_0^\pi e^{\eta \cos^2 \theta} h(\cos \theta) \cos \theta \sin^n \theta d\theta.$$

Using (98) and (52), this can be changed into

$$\tilde{\gamma}_1 = -\frac{\rho}{2\eta(n-1)} \frac{C_n}{Z_\eta} \int_0^\pi e^{\eta \cos^2 \theta} g(\theta) \frac{d\tilde{U}_0}{d\theta} \sin^{n-2} \theta d\theta$$

But from (45), we have $\frac{Z_\eta}{C_n} = \int_0^\pi e^{\eta \cos^2 \theta} \sin^{n-2} \theta d\theta$, which leads to the second equation in Formula (132). \square

REFERENCES

- [1] J. M. Ball, Axisymmetry of critical points for the Onsager functional, *Phil. Trans. R. Soc. A.*, **379** (2021), Paper No. 20200110, 13 pp.
- [2] J. M. Ball, [Mathematics and liquid crystals](#), *Molecular Crystals and Liquid Crystals*, **647** (2017), 1–27.
- [3] J. M. Ball, E. Feireisl and F. Otto, *Mathematical Thermodynamics of Complex Fluids*, Lecture notes in Mathematics 2200, Springer, 2017.
- [4] C. Bardos, F. Golse and C. D. Levermore, [Fluid dynamic limits of kinetic equations II convergence proofs for the Boltzmann equation](#), *Commun. Pure Appl. Math.*, **46** (1993), 667–753.
- [5] C. Bardos, F. Golse and D. Levermore, [Fluid dynamic limits of kinetic equations. I. Formal derivations](#), *J. Stat. Phys.*, **63** (1991), 323–344.
- [6] L. Boltzmann, [Weitere Studien über das Wärmegleichgewicht unter Gas-molekülen](#), *Sitzungsberichte Akad. Wiss., Vienna, part II*, **66** (1872), 275–370.
- [7] R. E. Caflisch, [The fluid dynamical limit of the nonlinear Boltzmann equation](#), *Commun. Pure Appl. Math.*, **33** (1980), 651–666.
- [8] M. C. Calderer, D. Golovaty, F.-H. Lin and C. Liu, [Time evolution of nematic liquid crystals with variable degree of orientation](#), *SIAM J. Math. Anal.*, **33** (2002), 1033–1047.
- [9] M. C. Calderer and C. Liu, [Liquid crystal flow: Dynamic and static configurations](#), *SIAM J. Appl. Math.*, **60** (2000), 1925–1949.
- [10] C. Cercinani, R. Illner and M. Pulvirenti, *The Mathematical Theory of Dilute Gases*, Springer, 2013.
- [11] S. Chapman, The kinetic theory of simple and composite gases: Viscosity, thermal conduction and diffusion, *Proc. Roy. Soc., (London)* **A93** (1916/17), 1–20.
- [12] B. Charbonneau, P. Charbonneau, Y. Jin, G. Parisi and F. Zamponi, [Dimensional dependence of the Stokes–Einstein relation and its violation](#), *The Journal of Chemical Physics*, **139** (2013), 164502.
- [13] X. Chen and J.-G. Liu, [Global weak entropy solution to Doi-Saintillan-Shelley model for active and passive rod-like and ellipsoidal particle suspensions](#), *J. Diff. Eqs.*, **254** (2013), 2764–2802.
- [14] P. Constantin, I. G. Kevrekidis and E. S. Titi, [Asymptotic states of a Smoluchowski equation](#), *Arch. Ration. Mech. Anal.*, **174** (2004), 365–384.
- [15] P. Constantin and J. Vukadinovic, [Note on the number of steady states for a two-dimensional Smoluchowski equation](#), *Nonlinearity*, **18** (2005), 441–443.
- [16] P.-G. de Gennes and J. Prost, *The Physics of Liquid Crystals*, Oxford Univ. Press, 1993.

- [17] P. Degond, A. Diez, A. Frouvelle and S. Merino-Aceituno, [Phase transitions and macroscopic limits in a BGK model of body-attitude coordination](#), *J. Nonlinear Sci.*, **30** (2020), 2671–2736.
- [18] P. Degond, G. Dimarco, T. B. N. Mac and N. Wang, [Macroscopic models of collective motion with repulsion](#), *Commun. Math. Sci.*, **13** (2015), 1615–1638.
- [19] P. Degond, A. Frouvelle, S. Merino-Aceituno and A. Trescases, [Quaternions in collective dynamics](#), *Multiscale Model. Simul.*, **16** (2018), 28–77.
- [20] P. Degond, J.-G. Liu, S. Motsch and V. Panferov, [Hydrodynamic models of self-organized dynamics: Derivation and existence theory](#), *Methods Appl. Anal.*, **20** (2013), 89–114.
- [21] P. Degond and S. Merino-Aceituno, [Nematic alignment of self-propelled particles: From particle to macroscopic dynamics](#), *Math. Models Methods Appl. Sci.*, **30** (2020), 1935–1986.
- [22] P. Degond, S. Merino-Aceituno, F. Vergnet and H. Yu, [Coupled self-organized hydrodynamics and Stokes models for suspensions of active particles](#), *J. Math. Fluid Mech.*, **21** (2019), Paper No. 6, 36 pp.
- [23] P. Degond and S. Motsch, [Continuum limit of self-driven particles with orientation interaction](#), *Math. Models Methods Appl. Sci.*, **18** (2008), 1193–1215.
- [24] M. Doi and S. F. Edwards, *The Theory of Polymer Dynamics*, Oxford Univ. Press, 1986.
- [25] W. E and P. Zhang, [A molecular kinetic theory of inhomogeneous liquid crystal flow and the small Deborah number limit](#), *Methods Appl. Anal.*, **13** (2006), 181–198.
- [26] D. Enskog, *Kinetische Theorie der Vorgänge in Mässig Verdünntent Gasen, 1*, in Allgemeiner Teil, Almqvist & Wiksell, Uppsala, 1917.
- [27] J. L. Ericksen, [Liquid crystals with variable degree of orientation](#), *Arch. Ration. Mech. Anal.*, **113** (1990), 97–120.
- [28] I. Fatkullin and V. Slastikov, [A note on the Onsager model of nematic phase transitions](#), *Commun. Math. Sci.*, **3** (2005), 21–26.
- [29] I. Fatkullin and V. Slastikov, [Critical points of the Onsager functional on a sphere](#), *Nonlinearity*, **18** (2005) 2565–2580.
- [30] A. Frouvelle, [A continuum model for alignment of self-propelled particles with anisotropy and density-dependent parameters](#), *Math. Models Methods Appl. Sci.*, **22** (2012), 1250011, 40 pp.
- [31] A. Frouvelle, [Body-attitude alignment: First order phase transition, link with rodlike polymers through quaternions, and stability](#), [arXiv:2011.14891](#), 2021.
- [32] Y. Giga and A. Novotný (eds.), *Handbook of Mathematical Analysis in Mechanics of Viscous Fluids*, Springer, 2018.
- [33] H. Grad, [On the kinetic theory of rarefied gases](#), *Commun. Pure Appl. Math.*, **2** (1949), 331–407.
- [34] D. Hilbert, [Begründung der kinetischen Gastheorie](#), *Mathematische Annalen*, **72** (1912), 562–577.
- [35] J. Huang, F. Lin and C. Wang, [Regularity and existence of global solutions to the Ericksen-Leslie system in \$\mathbb{R}^2\$](#) , *Comm. Math. Phys.*, **331** (2014), 805–850.
- [36] G. B. Jeffery, [The motion of ellipsoidal particles immersed in a viscous fluid](#), *Proceedings of the Royal Society A*, **102** (1922), 161–179.
- [37] N. Kuzuu and M. Doi, [Constitutive equation for nematic liquid crystals under weak velocity gradient derived from a molecular kinetic equation](#), *Journal of the Physical Society of Japan*, **52** (1983), 3486–3494.
- [38] F. M. Leslie, [Some constitutive equations for anisotropic fluids](#), *Quart. J. Mech. Appl. Math.*, **19** (1966), 357–370.
- [39] C. D. Levermore, [Moment closure hierarchies for kinetic theories](#), *J. Stat. Phys.*, **83** (1996), 1021–1065.
- [40] F.-H. Lin, [Nonlinear theory of defects in nematic liquid crystals; phase transition and flow phenomena](#), *Comm. Pure Appl. Math.*, **42** (1989), 789–814.
- [41] F.-H. Lin, [On nematic liquid crystals with variable degree of orientation](#), *Comm. Pure Appl. Math.*, **44** (1991), 453–468.
- [42] F.-H. Lin, J. Lin and C. Wang, [Liquid crystal flows in two dimensions](#), *Arch. Ration. Mech. Anal.*, **197** (2010), 297–336.
- [43] F.-H. Lin and C. Liu, [Nonparabolic dissipative systems modeling the flow of liquid crystals](#), *Comm. Pure Appl. Math.*, **48** (1995), 501–537.
- [44] F.-H. Lin and C. Liu, [Existence of solutions for the Ericksen-Leslie system](#), *Arch. Ration. Mech. Anal.*, **154** (2000), 135–156.

- [45] F.-H. Lin, C. Liu and P. Zhang, [On a micro-macro model for polymeric fluids near equilibrium](#), *Comm. Pure Appl. Math.*, **60** (2007), 838–866.
- [46] F.-H. Lin and T. Zhang, [Global small solutions to a complex fluid model in three dimensional](#), *Arch. Ration. Mech. Anal.*, **216** (2015), 905–920.
- [47] H. Liu, [Global orientation dynamics for liquid crystalline polymers](#), *Phys. D*, **228** (2007), 122–129.
- [48] H. Liu, H. Zhang and P. Zhang, [Axial symmetry and classification of stationary solutions of Doi-Onsager equation on the sphere with Maier-Saupe potential](#), *Commun. Math. Sci.*, **3** (2005), 201–218.
- [49] C. Luo, H. Zhang and P. Zhang, [The structure of equilibrium solutions of the one-dimensional Doi equation](#), *Nonlinearity*, **18** (2005), 379–389.
- [50] W. Maier and A. Saupe, A simple molecular statistical theory of the nematic crystalline-liquid phase, *I Z Naturf. A*, **14** (1959), 882–889.
- [51] J. C. Maxwell, On the dynamical theory of gases, *Philos. Trans. Roy. Soc. London*, **157** (1867), 49–88.
- [52] A. Mellet, [Fractional diffusion limit for collisional kinetic equations: A moments method](#), *Indiana Univ. Math. J.*, **59** (2010), 1333–1360.
- [53] L. Onsager, The effects of shape on the interaction of colloidal particles, *Ann. N. Y. Acad. Sci.*, **51** (1949), 627–659.
- [54] F. Otto and A. E. Tzavaras, [Continuity of velocity gradients in suspensions of rod-like molecules](#), *Comm. Math. Phys.*, **277** (2008), 729–758.
- [55] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen and O. Shochet, [Novel type of phase transition in a system of self-driven particles](#), *Phys. Rev. Lett.*, **75** (1995), 1226–1229.
- [56] H. Wang and P. J. Hoffman, [A unified view on the rotational symmetry of equilibria of nematic polymers, dipolar nematic polymers, and polymers in higher dimensional space](#), *Commun. Math. Sci.*, **6** (2008), 949–974.
- [57] Q. Wang, W. E, C. Liu and P.-W. Zhang, [Kinetic theory for flows of nonhomogeneous rodlike liquid crystalline polymers with a nonlocal intermolecular potential](#), *Phys. Rev. E*, **65** (2002), 051504.
- [58] W. Wang, L. Zhang and P. Zhang, [Modeling and computation of liquid crystals](#), *Acta Numer.*, **30** (2021), 765–851.
- [59] W. Wang, P. Zhang and Z. Zhang, [The small Deborah number limit of the Doi-Onsager equation to the Ericksen-Leslie equation](#), *Comm. Pure Appl. Math.*, **68** (2015), 1326–1398.
- [60] W. Wang, P. Zhang and Z. Zhang, [Well-posedness of the Ericksen-Leslie system](#), *Arch. Ration. Mech. Anal.*, **210** (2013), 837–855.
- [61] H. Zhang and P. Zhang, [Stable dynamic states at the nematic liquid crystals in weak shear flow](#), *Phys. D*, **232** (2007), 156–165.
- [62] H. Zhang and P. Zhang, [On the new multiscale rodlike model of polymeric fluids](#), *SIAM J. Math. Anal.*, **40** (2008), 1246–1271.
- [63] H. Zhou, H. Wang, M. G. Forest and Q. Wang, [A new proof on axisymmetric equilibria of a three-dimensional Smoluchowski equation](#), *Nonlinearity*, **18** (2005), 2815–2825.

Received June 2021; revised October 2021; early access January 2022.

E-mail address: pierre.degond@math.univ-toulouse.fr

E-mail address: frouvelle@ceremade.dauphine.fr, amic.frouvelle@math.univ-poitiers.fr

E-mail address: jliu@phy.duke.edu